

# Surgical and Ablative Procedures for Venous Insufficiency and Varicose Veins (for Indiana Only)

**Policy Number:** CS1171N.08  
**Effective Date:** December 1, 2023

[Instructions for Use](#)

Table of Contents	Page
<a href="#">Application</a> .....	1
<a href="#">Coverage Rationale</a> .....	1
<a href="#">Definitions</a> .....	2
<a href="#">Applicable Codes</a> .....	3
<a href="#">Description of Services</a> .....	5
<a href="#">Benefit Considerations</a> .....	6
<a href="#">Clinical Evidence</a> .....	6
<a href="#">U.S. Food and Drug Administration</a> .....	14
<a href="#">References</a> .....	14
<a href="#">Policy History/Revision Information</a> .....	17
<a href="#">Instructions for Use</a> .....	18

Related Policies
None

## Application

This Medical Policy only applies to the state of Indiana.

## Coverage Rationale

[See Benefit Considerations](#)

### Varicose Vein Ablative and Stripping Procedures

**Varicose Vein ablative and Stripping procedures are considered reconstructive, proven, and medically necessary in certain circumstances.** For medical necessity clinical coverage criteria, refer to the InterQual® CP: Procedures:

- Ablation, Endovenous, Varicose Vein
- Ligation/Excision, Varicose Vein, +/- Stripping

Click [here](#) to view the InterQual® criteria.

Refer to the [Coding Clarification](#) section. Adherence to American Medical Association (AMA) coding guidance is required when requesting coverage of Endovenous Ablation procedures. Note that only one primary code may be requested for the initial vein treated, and only one add-on code per extremity may be requested for any subsequent vein(s) treated.

### Ligation Procedures

**The following procedure is proven and medically necessary:**

- Ligation at the saphenofemoral junction, as a stand-alone procedure, when used to prevent the propagation of an active clot to the deep venous system in individuals with ascending Superficial Thrombophlebitis who fail or are intolerant of anticoagulation therapy

**The following procedure is proven and medically necessary in certain circumstances:**

- Ligation, subfascial, endoscopic surgery for treatment of perforating veins associated with chronic Venous Insufficiency. For medical necessity clinical coverage criteria, refer to the InterQual® CP: Procedures, Ligation, Subfascial, Endoscopic, Perforating Vein

Click [here](#) to view the InterQual® criteria.

**The following procedures are unproven and not medically necessary for treating Venous Reflux due to insufficient evidence of efficacy:**

- Ligation of the Great Saphenous Vein (GSV) at the saphenofemoral junction, as a stand-alone procedure
- Ligation of the Small Saphenous Vein (SSV) at the saphenopopliteal junction, as a stand-alone procedure
- Ligation at the saphenofemoral junction, as an adjunct to radiofrequency ablation or endovenous laser ablation of the main saphenous veins

## Ambulatory Phlebectomy

**Ambulatory phlebectomy for treating Varicose Veins is proven and medically necessary in certain circumstances.** For medical necessity clinical coverage criteria, refer to the InterQual® CP: Procedures, Ambulatory Phlebectomy, Varicose Vein for:

- Hook Phlebectomy
- Microphlebectomy
- Mini Phlebectomy
- Stab Avulsion
- Stab Phlebectomy

Click [here](#) to view the InterQual® criteria.

## Other Procedures

**Sclerotherapy is proven and medically necessary in certain circumstances.** For medical necessity clinical coverage criteria, refer to the InterQual® CP: Procedures, Sclerotherapy, Varicose Vein.

Click [here](#) to view the InterQual® criteria.

**The following procedures are unproven and not medically necessary for treating Venous Reflux due to insufficient evidence of efficacy:**

- Endovascular embolization of Varicose Veins using cyanoacrylate-based adhesive
- Endovenous mechanochemical ablation (MOCA) of Varicose Veins
- Porcine bioprosthetic valve (e.g., VenoValve) implantation into the femoral vein for treatment of deep vein reflux associated with chronic Venous Insufficiency

## Definitions

Check the definitions within the federal, state, and contractual requirements that supersede the definitions below.

**Endovenous Ablation:** A minimally invasive procedure that uses heat generated by radiofrequency (RF) or laser energy to seal off damaged veins.

**Great Saphenous Vein (GSV):** The GSV originates from the dorsal arch of the foot and progresses medially and proximally along the distal extremity to join the common femoral vein.

**Ligation:** Tying off a vein.

**Reticular Vein:** Reticular Veins are dilated dermal veins less than 4 mm in diameter that communicate with either or both Telangiectasia and saphenous tributaries.

**Small Saphenous Vein:** Superficial vein of the calf.

**Spider Vein:** Spider Veins/Telangiectasia are the permanent dilation of preexisting small blood vessels, generally up to 1mm in size.

**Stripping:** Surgical removal of superficial veins.

**Superficial Thrombophlebitis:** Inflammation of a vein due to a blood clot in a vein just below the skin's surface.

**Telangiectasia:** Refer to [Spider Vein](#).

**Varicose Veins:** Abnormally enlarged veins that are frequently visible under the surface of the skin; often appear blue, bulging and twisted.

**Venous Reflux/Insufficiency:** Venous Reflux is reversed blood flow in the veins (away from the heart). Abnormal (pathological reflux) is defined as reverse flow that lasts beyond a specified period of time as measured by Doppler ultrasound. Normal (physiological reflux) is defined as reverse flow that lasts less than a specified period of time as measured by Doppler ultrasound. Abnormal (pathological reflux) times exceed different thresholds depending on the system of veins:

- Deep veins: 1 sec
- Superficial veins: 0.5 sec
- Perforator veins: 0.35 sec

## Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

### Coding Clarifications:

- According to the American Medical Association (AMA), CPT code 37241 is specific to venous embolization/occlusion and excludes lower extremity venous incompetency. Coding instructions state that 37241 should not be used to request treatment of incompetent extremity veins. For sclerosis of veins or endovenous ablation of incompetent extremity veins, refer to 36468-36479 (CPT Assistant, 2014).
- Adherence to AMA coding guidance is required when requesting endovenous ablation procedures.

Per AMA coding guidance, the initial incompetent vein treated (e.g., [36475](#)) may only be requested once per extremity. For endovenous ablation, treatment of subsequent incompetent veins in the same extremity as the initial vein treated (e.g., [36476](#)), only one add-on code per extremity may be requested, regardless of the number of additional vein(s) treated (CPT Assistant, November 2016).

Therefore, only one primary code may be requested for the initial vein treated, and only one add-on code per extremity may be requested for any subsequent vein(s) treated.

CPT Code	Description
*0744T	Insertion of bioprosthetic valve, open, femoral vein, including duplex ultrasound imaging guidance, when performed, including autogenous or nonautogenous patch graft (e.g., polyester, ePTFE, bovine pericardium), when performed
*36465	Injection of non-compounded foam sclerosant with ultrasound compression maneuvers to guide dispersion of the injectate, inclusive of all imaging guidance and monitoring; single incompetent extremity truncal vein (e.g., great saphenous vein, accessory saphenous vein)

CPT Code	Description
*36466	Injection of non-compounded foam sclerosant with ultrasound compression maneuvers to guide dispersion of the injectate, inclusive of all imaging guidance and monitoring; multiple incompetent truncal veins (e.g., great saphenous vein, accessory saphenous vein), same leg
*36468	Injection(s) of sclerosant for spider veins (telangiectasia), limb or trunk
*36470	Injection of sclerosant; single incompetent vein (other than telangiectasia)
*36471	Injection of sclerosant; multiple incompetent veins (other than telangiectasia), same leg
36473	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, mechanochemical; first vein treated
*36474	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, mechanochemical; subsequent vein(s) treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure)
36475	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, radiofrequency; first vein treated
*36476	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, radiofrequency; subsequent vein(s) treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure)
36478	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, laser; first vein treated
*36479	Endovenous ablation therapy of incompetent vein, extremity, inclusive of all imaging guidance and monitoring, percutaneous, laser; subsequent vein(s) treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure)
*36482	Endovenous ablation therapy of incompetent vein, extremity, by transcatheter delivery of a chemical adhesive (e.g., cyanoacrylate) remote from the access site, inclusive of all imaging guidance and monitoring, percutaneous; first vein treated
*36483	Endovenous ablation therapy of incompetent vein, extremity, by transcatheter delivery of a chemical adhesive (e.g., cyanoacrylate) remote from the access site, inclusive of all imaging guidance and monitoring, percutaneous; subsequent vein(s) treated in a single extremity, each through separate access sites (List separately in addition to code for primary procedure)
*37500	Vascular endoscopy, surgical, with ligation of perforator veins, subfascial (SEPS)
37700	Ligation and division of long saphenous vein at saphenofemoral junction, or distal interruptions
37718	Ligation, division, and stripping, short saphenous vein
37722	Ligation, division, and stripping, long (greater) saphenous veins from saphenofemoral junction to knee or below
*37735	Ligation and division and complete stripping of long or short saphenous veins with radical excision of ulcer and skin graft and/or interruption of communicating veins of lower leg, with excision of deep fascia
*37765	Stab phlebectomy of varicose veins, 1 extremity; 10-20 stab incisions
*37766	Stab phlebectomy of varicose veins, 1 extremity; more than 20 incisions
37780	Ligation and division of short saphenous vein at saphenopopliteal junction (separate procedure)
*37785	Ligation, division, and/or excision of varicose vein cluster(s), 1 leg
*37799	Unlisted procedure, vascular surgery

*CPT® is a registered trademark of the American Medical Association*

**Note:** Codes labeled with an asterisk (\*) are not managed for medical necessity review for the state of Indiana at the time this policy became effective. Refer to the most up to date prior authorization list for Indiana at [Prior Authorization and Notification: UnitedHealthcare Community Plan of Indiana](#).

## Description of Services

Varicose Veins are enlarged veins that are swollen and raised above the surface of the skin. They can be dark purple or blue and look twisted and bulging. Varicose Veins are commonly found on the backs of the calves or on the inside of the leg. Veins have one-way valves that help keep blood flowing towards the heart. When the valves become weak or damaged and do not close properly, blood can back up and pool in the veins causing them to get larger. The resulting condition is known as Venous Insufficiency or Venous Reflux. Varicose Veins may lead to complications such as pain, blood clots or skin ulcers.

Duplex ultrasound is considered the “gold standard” for diagnosis of superficial venous incompetence. The CEAP (clinical, etiology, anatomy, pathophysiology) classification system is used to describe the degree of varicosity. The “C” part of CEAP classification is more useful and practical in rating the severity of Varicose Veins:

- C0: No visible or palpable signs of venous disease
- C1: Telangiectasias (Spider Veins) or Reticular Veins
- C2: Varicose Veins (diameter of vein is > 3mm)
- C3: Edema
- C4a: Pigmentation and eczema
- C4b: Lipodermatosclerosis and atrophie blanche
- C5: Healed venous ulcer
- C6: Active venous ulcer

(Lurie et al. (American Venous Forum [AVF]), 2020)

Venous clinical severity scoring has been used to measure clinical improvement after treatment of Varicose Veins. Other venous severity scoring methods include Venous Severity Score, Venous Clinical Severity Score, Venous Segmental Disease Score (Lurie et al. (AVF), 2020).

Preoperative venous duplex ultrasound is used to evaluate patients for Venous Insufficiency symptoms or suspected DVT; it can provide a road map of vein anatomy similar to contrast venography, as well as essential hemodynamic information about the presence of proximal obstruction, vein valve function, and Venous Reflux (Lin et al., 2015).

Varicose Veins are treated with lifestyle changes and medical procedures done either to remove the veins or to close them. Endovenous Ablation therapy uses lasers or radiofrequency energy to create heat to close off a Varicose Vein. Vein Stripping and Ligation involves tying shut and removing the veins through small cuts in the skin (National Heart, Lung and Blood Institute [NHLBI], 2014).

Endomechanical ablation uses a specialized, rotating catheter (e.g., ClariVein) to close off a Varicose Vein by damaging the vessel lining prior to injecting a sclerosing agent. This technique is also referred to as mechanochemical ablation (MOCA), mechanico-chemical endovenous ablation (MCEA) and mechanically enhanced endovenous chemical ablation (MEECA).

Endovascular embolization using cyanoacrylate-based adhesive (e.g., VenaSeal™ Closure System) is a minimally invasive, non-thermal and non-sclerosant procedure that does not require tumescent anesthesia. The medical adhesive is used to close the lower extremity superficial truncal veins, such as the Great Saphenous Vein, in individuals with symptomatic Venous Reflux disease.

Endovascular embolization using endovenous foam sclerotherapy with polidocanol endovenous microfoam (PEM) (e.g., Varithena™ [Provensis Ltd.]), is a prescribed proprietary canister that generates a sterile, uniform, stable, low-nitrogen polidocanol 1% microfoam sclerosant intended for ultrasound-guided intravenous (IV) injection for treating venous incompetence and varicosities (Hayes, 2022). The aim of ultrasound-guided foam sclerotherapy for Varicose Veins is to damage the endothelial surface of the vein causing scarring and leading to blockage of the treated Varicose Veins. Sclerosant, in the form of a foam, is intended to have good surface area contact with the vein walls (National Institute of Health and Care Excellence [NICE], 2013).

# Benefit Considerations

## Coverage Limitations and Exclusions

- The following procedures are excluded from coverage: Any procedure that does not meet the criteria in the [Coverage Rationale](#) section. Treatments for Spider Veins and/or Telangiectasias are considered to be cosmetic and therefore excluded from coverage.
- Endovenous Ablation (radiofrequency and/or laser) of either reticular or telangiectatic veins is not reconstructive and not medically necessary and therefore excluded from coverage.

## Clinical Evidence

### Endovenous Mechanochemical Ablation

Evidence in peer review literature evaluating endovenous mechanochemical ablation (MOCA) for the treatment of venous insufficiency and varicose veins is limited. Future robust RCTs are warranted along with long-term outcomes to establish the safety and efficacy of this procedure.

A Hayes Health Technology Assessment states MOCA with the ClariVein infusion catheter appears safe and effective over the short-term but the low-quality body of evidence does not allow conclusions to be drawn regarding the long-term durability of the procedure. The report states that MOCA resulted in slightly poorer technical outcomes and higher rates of recanalization than thermal ablation and surgical procedures. The report recommends future well-designed trials with larger sample sizes that compare MOCA using the ClariVein infusion catheter with clinical alternatives with a long-term follow-up (Hayes, 2022).

Mohamed et al. (2021) conducted a single-center RCT to compare the technical, clinical and QoL outcomes after EVLA and MOCA. One hundred fifty patients with symptomatic, unilateral, single-axis superficial venous incompetence (SVI) were randomized equally to either EVLA or MOCA, both with concomitant phlebectomy when necessary. Primary outcomes were intraprocedural axial ablation pain scores and anatomical occlusion at one year. Secondary outcomes included postprocedural pain, VCSS, QOL (Aberdeen Varicose Veins Questionnaire [AVVQ] and EuroQol 5-domain utility index), patient satisfaction and complication rates. Both groups reported low intraprocedural pain scores; on a 100 mm visual analog scale, pain during axial EVLA was 22 (9-44) compared to 15 (9-29) during MOCA. At 1-year, duplex derived anatomical occlusion rates after EVLA were 63/69 (91%) compared to 53/69 (77%) in the MOCA group ( $p = 0.02$ ). Both groups experienced improvement in VCSS and AVVQ after treatment, without a significant difference between groups. Median VCSS improved from six (5-8) to zero (0-1) at one year. Median AVVQ improved from 13.8 (10.0-17.7) to 2.0 (0.0-4.9). One patient in the MOCA group experienced DVT. The authors concluded EVLA resulted in a higher technical success rate compared to MOCA but clinically, both treatments were highly efficacious in treating SVI. Patients improved in terms of symptoms, disease severity and QoL. Both procedures resulted in low procedural pain with a short recovery time. EVLA had higher axial occlusion rates. The authors noted that higher recanalization rates after MOCA may lead to higher rates of recurrence and long-term follow-up is needed. Long-term follow up at five and 10 years is planned for this study. Limitations include short term follow up and single-center recruitment.

In an updated Cochrane review, Whing et al. (2021) compared interventions for treating varicosities of the GSV. The review included 24 RCTs with 5,135 participants who underwent EVLA, RFA, EVSA, UGFS, cyanoacrylate glue, MOCA, or high ligation and stripping. The authors found there was no clear difference in technical success or recurrence between RFA compared to MOCA, however, long-term data were not available, and the confidence intervals of the combined data were broad, making these findings largely inconclusive. Additionally, the authors noted all the trials had some risk of bias concerns. The authors determined there were a relatively small number of studies for comparison and differences in outcome definitions and time points reported limited their conclusions. Future studies which provide more evidence on the breadth of treatments are recommended by the authors. Bootun et al. (2016), Lane et al. (2017), Holewijn et al. (2019), Vähäaho et al. (2019), which were previously cited in this policy, are included in this review.

Kim et al. (2017) evaluated in a case series whether early efficacy in endovenous MOCA is maintained at 24 months. Patients with reflux in the GSV involving the sapheno-femoral junction and no previous venous interventions were included. The occlusion rate of treated veins was assessed with duplex ultrasound. Patient clinical improvement was assessed by CEAP class and VCSS. Of the initial 126 patients, there were 65 patients with 24-month follow-up. Of these 65 patients, 70% were female, with a mean age of  $70 \pm 14$  years and an average BMI of  $30.5 \pm 6$ . The mean GSV diameter in the upper thigh was 7.6 mm and

the mean treatment length was 39 cm. Adjunctive treatment of the varicosities was performed in 14% of patients during the procedure. Closure rates were 100% at one week, 98% at three months, 95% at 12 months, and 92% at 24 months. There was one patient with complete and four with partial recanalization ranging from 7 to 12 cm (mean length 9 cm). There was significant improvement in CEAP and VCSS ( $p < .001$ ) for all time intervals. Early high occlusion rate with MOCA is associated with significant clinical improvement, which was maintained at 24 months. According to the authors, this finding is suggestive of a good option for the treatment of GSV incompetence. Longer-term outcomes are needed to evaluate MOCA's efficacy. The study is limited by lack of comparison group and large loss to follow-up.

Vos et al. (2017) conducted a systematic review and meta-analysis to evaluate the efficacy of MOCA and cyanoacrylate vein ablation (CAVA) for GSV incompetence. Eligible articles were prospective studies that included patients treated for GSV incompetence and described the primary outcome. Exclusion criteria were full text not available, case reports, retrospective studies, small series ( $n < 10$ ), reviews, abstracts, animal studies, studies of SSV incompetence, and recurrent GSV incompetence. Primary outcome was anatomic success. Secondary outcomes were initial technical success, VCSS, AVVQ score, and complications. Fifteen articles met the inclusion criteria. Pooled anatomic success for MOCA and CAVA was 94.7% and 94.8% at six months and 94.1% and 89.0% at one year, respectively. VCSS and AVVQ score significantly improved after treatment with MOCA and CAVA. The authors conclude that both of these non-thermal techniques are promising that could serve as alternatives for thermal ablation techniques. However, to determine their exact role in clinical practice, high-quality RCTs comparing these novel modalities with well-established techniques are required. This study is limited by inclusion or mostly uncontrolled studies to assess the efficacy and safety of MOCA. Elias and Raines (2012) and Bishawi et al. (2014), which were previously cited in this policy, are included in this meta-analysis.

Witte et al. (2017a) conducted a systematic review and meta-analysis of MOCA of saphenous veins using the ClariVein to report on the anatomical, technical, and clinical success. The literature search identified 759 records, of which 13 were included, describing 10 unique cohorts. A total of 1,521 veins (1,267 GSV and 254 SSV) were included, with cohort sizes ranging from 30 to 570 veins. The pooled anatomical success rate after short-term follow up was 92% (95% CI 90-94%) ( $n = 1,314$  veins). After 6 and 12 months these numbers were 92% (95% CI 88-95%) ( $n = 284$ ) and 91% (95% CI 86-94%) ( $n = 228$ ), respectively. The long-term anatomical success rates at two and three years were 91% (95% CI 85-95%) ( $n = 136$ ) and 87% (95% CI 75-94%) ( $n = 48$ ), respectively. Major complications and especially nerve injury were very rare ( $\leq 0.2\%$ ). All studies were of moderate or good quality using the methodological index for non-randomized studies (MINORS) scoring scale. The authors concluded that MOCA using the ClariVein in combination with liquid sclerosant is associated with an anatomical success rate ranging from 87% to 92% and good clinical success. However, they reported that no RCTs are available studying the anatomical success after MOCA compared to the endothermal ablation.

Witte et al. (2017b) reported midterm results of MOCA for treating GSV insufficiency. In a 1-year period, 85 consecutive patients undergoing MOCA with polidocanol in 104 limbs were enrolled in a prospective registry. The patients were evaluated at baseline and during follow-up (four weeks and one, two, and three years) using duplex ultrasound, the CEAP classification, the VCSS, the RAND Short Form 36-Item Health Survey (RAND-SF36), and the AVVQ. Primary outcome measures were clinical and anatomic success. Secondary outcome measures included general and disease specific QoL and re-interventions. After a median follow-up of 36 months (interquartile range 12.5, 46.3), recanalization occurred in 15 (15%) of 102 successfully treated vein segments. Anatomic success was 92%, 90%, and 87% after one, two, and three years, respectively. The VCSS improved at all time intervals compared to the preprocedure median. The clinical success at three years was 83%. The AVVQ and RAND-SF36 scores showed an improvement at all time intervals compared to baseline values. Between 12 and 36 months, however, a significant deterioration was observed in VCSS, which was accompanied by worsening of disease specific and general QOL. Although the authors concluded that MOCA demonstrated to be an effective treatment modality for GSV insufficiency at midterm follow-up, clinical results seemed to drop over time. Additionally, these findings are limited by lack of comparison group undergoing a different treatment.

Vun et al. (2015) assessed the efficacy of the ClariVein system for the treatment of superficial vein incompetence. Fifty-one GSVs and six SSVs were treated. Duplex showed a technical success rate of 91%. Comparison with 50 RFA and 40 EVLA procedures showed procedure times were significantly less for ClariVein than for either RFA or EVLA. Median pain scores were significantly lower for ClariVein than for RFA and EVLA. No major complications or deep vein thromboses were reported. Study limitations included small sample size, lack of randomization and short-term follow-up. Further data on long-term clinical outcomes is needed.

In a prospective case series, Boersma et al. (2013) evaluated the feasibility, safety and 1-year results of MOCA of SSV insufficiency. Fifty consecutive patients were treated using the ClariVein device and polidocanol. At the 6-week assessment, all treated veins were occluded. One-year follow-up showed a 94% anatomic success rate and no major complications. The authors concluded that MOCA is a safe, feasible and efficacious technique for treating SSV insufficiency. This study is limited by lack of control group, small sample size and short-term follow-up.

In a prospective comparison study, van Eekeren et al. (2013) evaluated postoperative pain and QOL after RFA and MOCA for GSV incompetence. Sixty-eight patients with unilateral GSV incompetence were included. Patients treated with MOCA reported significantly less postoperative pain than patients treated with RFA during the first 14 days after treatment. The lower postoperative pain score was associated with a significantly earlier return to normal activities and work. At 6 weeks, patients in both groups perceived an improved change in health status and an improved disease specific QOL. This study is limited by lack of randomization, small sample size and short-term follow-up.

In a pilot study, van Eekeren et al. (2011) evaluated the feasibility and safety of endovenous MOCA for the treatment of GSV incompetence. Thirty limbs in 25 patients (18 women; mean age 52 years) with GSV incompetence were treated with the ClariVein® device. Initial technical success, complications, patient satisfaction and classification by VCSS were assessed 6 weeks after the treatment. Initial technical success of MOCA was 100%. There were no major adverse events. Duplex ultrasonography at six weeks showed 26 (87%) of 30 veins were completely occluded. Three veins showed partial recanalization in the proximal and distal GSV. One patient had full segment recanalization and was successfully retreated. The VCSS significantly improved at six weeks. Patient satisfaction was high, with a median satisfaction of 8.8 on a 0-10 scale. The authors concluded that endovenous MOCA is feasible and safe in the treatment of GSV incompetence. Larger studies with a prolonged follow-up are indicated to prove the efficacy of this technique. This study is limited by lack of comparison group undergoing a different treatment approach.

## Endovascular Embolization with Cyanoacrylate-Based Adhesive

Quality evidence in peer review literature evaluating endovascular embolization with cyanoacrylate-based adhesive for the treatment of venous insufficiency and varicose veins is limited. Future robust RCTs are warranted along with long-term outcomes to establish the safety and efficacy of this procedure. An ongoing RCT may provide more definitive findings about this technology (NCT03820947).

A 2022 Hayes Health Technology Assessment evaluated nine clinical studies on the efficacy and safety of cyanoacrylate embolization with the VenaSeal Closure System. The evidence included three RCTs and six retrospective comparative studies. The conclusion states that a low-quality body of evidence suggests VenaSeal has a high level of successful venous closure for at least one year that may result in reduced symptom severity and improved QoL. Efficacy and safety may be comparable to RFA, EVLA, and MOCA; however, substantial uncertainty remains regarding its effectiveness due to the lack of well-designed comparative studies and limited follow-up beyond one year. The authors overall conclusion is that cyanoacrylate embolization with the VenaSeal Closure System has potential but unproven benefits.

Amshar et al. (2022) conducted a systematic review and meta-analysis to evaluate the efficacy, intervention time, and safety of cyanoacrylate embolization (CAE) in comparison to EVLA in treatment of saphenous vein insufficiency. Efficacy was determined by venous closure rate one-year post-intervention and VCSS one-year post-intervention. Safety was determined by rates of periprocedural pain, skin pigmentation, nerve damage, phlebitis, DVT and ecchymosis. Two randomized-controlled trials and three cohort studies were included in this review. The total number of individuals was 1,432 (710 CAE and 722 EVLA). Venous closure rates and VCSS did not differ significantly between CAE group and EVLA group. Pooled data showed that CAE group was associated with less periprocedural pain score ( $p < 0.001$ ), lower skin pigmentation rates (0.60% vs. 4.46%;  $p = 0.008$ ), and lower nerve damage rates (0% vs. 3.94%;  $p = 0.007$ ). Rates of phlebitis, DVT, and ecchymosis did not differ significantly between the two groups. In addition, intervention time was significantly faster in CAE group compared to EVLA group ( $p < 0.001$ ). The authors concluded CAE was not inferior to EVLA in terms of efficacy and CAE showed less adverse effects occurrence rates of periprocedural pain, skin pigmentation, and nerve damage complications. Additionally, intervention time is stated to be faster with CAE compared to EVLA. The authors note that future RCTs with larger sample sizes and longer post-procedural follow-up time are needed. Additionally, efficacy outcomes were limited to one year and longer-term outcome data would provide additional evidence of efficacy. Bozkurt and Yilmaz (2016), and Eroglu and Yasim (2018) which were previously cited in this policy, are included in this review. Currently, the VariClose Vein Sealing System (Biolas, FG Grup, Turkey) is under research in countries other than the United States and has neither been approved nor cleared for marketing by the FDA.



An ECRI clinical evidence assessment (2021) suggests that VenaSeal is safe and as effective as RFA for treating varicose veins in patients with venous reflux disease. However, how well VenaSeal works compared with other treatment modalities cannot be determined because the SR assessed too few patients for each comparison and no studies in the SR performed head-to-head comparisons. The report determined the evidence was somewhat favorable but RCTs are needed to compare VenaSeal with other treatment modalities. Limitations of the reviewed studies include risk for lack of blinding, single-center focus, and lack of randomization.

Joh et al. (2021) conducted an open-label multicenter, prospective, RCT that compare the clinical outcomes of cyanoacrylate closure (CAC) and surgical stripping (SS) for the treatment of incompetent great saphenous veins. One hundred and twenty-six patients were randomized into two groups (63 with CAC and 63 with SS). Target vein occlusion was assessed on the third day and one, three, six, and 12 months postoperatively using duplex ultrasound. The primary endpoint of the study was to evaluate complete closure of the target vein at three months. Ecchymosis grades, VCSS, AVVQ score, and pain were also assessed as secondary outcomes. Postoperative pain scores were significantly better in the CAC group than in the SS group. In addition, the mean ecchymosis grade was  $0.3 \pm 0.5$  in the CAC group and  $1.1 \pm 1.1$  in the SS group ( $p < .001$ ). The VCSS and QoL had improved equally in both groups. Most complications were minor (nine events in CAC group and 20 events in SS group) with one major complication occurring in a patient who had undergone the SS procedure. Complete occlusion of the target vein at three months was achieved by both procedures. Postoperative pain and ecchymosis grades were significantly lower in the CAC group. The authors concluded that CAC has a high success rate with few complications. Limitations noted by the authors include lack of information on patient return to work and daily activities, pain scores during the procedure and immediately after the procedure were not obtained, the 2X2 factorial design with 1:1 randomization, could contribute to differences in gender distribution and VCSSs in the two groups and concomitant phlebectomy could have also influenced the occurrence of complications. Additionally, lack of masking could have introduced a bias in the findings.

The VenaSeal Saphen Closure System Pivotal Study (VeClose) is a multi-center RCT that compared cyanoacrylate closure (CAC) to RFA for the treatment of incompetent great saphenous veins. In this trial, 222 subjects with symptomatic GSV incompetence were randomly assigned to receive either CAC ( $n = 108$ ) with the VenaSeal Saphen Closure System or RFA ( $n = 114$ ). The primary endpoint was closure of the target vein at month three, as assessed by duplex ultrasound. To determine non-inferiority of CAE to RFA, the investigators used a predetermined margin of 10%. Secondary endpoints included subject-rated pain experienced during the procedure (i.e., pain experienced after vein access but before all treatment/access catheters were removed), investigator-rated ecchymosis at day three, adverse events, and details of adjunctive procedures. Patient follow-up visits were on day three and at months one, three, six, 12, 24, and 36. For the extension study, patients who were successfully contacted and were interested in participation provided written informed consent for the 60-month follow-up visit. Assessment tools included the VCSS, AVVQ and EuroQoL-Five Dimension (EQ-5D) quality of life survey. This trial has generated multiple publications that reported outcomes with various follow-up periods e.g., three months (Morrison, 2015), 12 months (Morrison, 2017) 24 months (Gibson, 2018a), 36 months (Morrison, 2019), and 60 months (Morrison, 2020), as well as a publication with results of a roll-in phase analysis, which included 20 additional patients treated with CAC (Kolluri, 2016). Design limitations of this study and the resulting publications included lack of blinding of the subjects or assessors to the intervention. Furthermore, the primary endpoint of the study was complete closure of the target vein at three months after index treatment, thus the study may not have been powered to detect clinically significant differences between treatments groups for important outcomes and at different times of follow-up. These studies were also included in the Hayes report (2022). The individual studies are listed below:

- Morrison et al. (2015) reported 3-month outcomes from the VeClose trial. No adjunctive procedures such as phlebectomy and UGFS were allowed until after the month three visit. The closure rates were 99% for VenaSeal and 96% for RFA. Pain experienced during the procedure was reported as mild and was similar between treatment groups. Good safety profiles were reported with both treatments. The authors concluded that cyanoacrylate ablation did not require tumescent anesthesia, was associated with less post procedure ecchymosis, and was noninferior to RFA for the treatment of incompetent GSVs at month three after the procedure.
- Morrison et al. (2017) reported 12-month outcomes from the VeClose trial. Of 222 randomized patients, a 12-month follow-up was obtained for 192 (95 CAC and 97 RFA; total follow-up rate, 86.5%). The complete occlusion rate was nearly identical in both groups (97.2% in the CAC group and 97.0% in the RFA group). Twelve-month freedom from recanalization was similar in the CAC and RFA groups, although there was a trend toward greater freedom from recanalization in the CAC group ( $p = .08$ ). The authors reported that patient symptoms and QOL improved equally in both groups.
- Twenty-four-month outcomes from the VeClose trial were reported by Gibson et al (2018a). One hundred and seventy-one patients completed the 24-month follow-up, which included 87 from the CAC group and 84 from the RFA group. The 24-month GSV closure rate was 95.3% in the CAC group and 94.0% in the RFA group. Symptoms and QOL improved similarly

in both groups. No clinically significant device- or procedure-related late adverse events were reported. The authors concluded that both CAC and RFA were effective in closure of the target GSV, resulting in similar and significant improvements in the patient's QoL through 24 months.

- One hundred and forty-six patients completed the 36-month follow-up to the VeClose trial, which included 72 patients from the CAC group and 74 patients from the RFA group, with outcomes reported by Morrison et al. (2019). The 36-month GSV closure rate was 94.4% for the CAC group and 91.9% for the RFA group. Stable improvement in symptoms and QOL was observed in both groups. Adverse event rates between the 24- and 36-month visits were similar between the groups as were serious adverse events which were infrequent and judged unrelated to either the device or the procedure in both groups. The authors surmised the results of this trial continue to demonstrate the safety and efficacy of CAC for the treatment of GSV incompetence with vein closure rate at 36 months similar to that of RFA. The findings are limited by the loss to follow up (34%), which could have introduced biases in the findings.
- Morrison et al. (2020) reported 60-month outcomes from the VeClose trial with a total of 89 patients in the original study completing the 60-month visit. Of those, 47 patients were from the CAC group, 33 patients were from the RFA group, and nine patients were from the roll-in CAC group. No new recanalization events were observed between 36 and 60 months of follow-up. Kaplan-Meier estimates for freedom from recanalization in the randomized CAC and RFA groups were 91.4% and 85.2%, respectively. Both groups demonstrated sustained improvements in EuroQol-5 Dimension (EQ-5D) and QOL. Whereas patients assigned to C0 or C1 clinical class were excluded from the original study, more than half of all returning patients (64% [57/89]) were now assigned to C0 or C1, suggesting an improved clinical class from baseline. Furthermore, 41.1% of returning CAC patients and 39.4% of returning RFA patients at least two CEAP clinical classes lower than at baseline. The authors concluded that CAC and RFA were effective in achieving complete target vein closure of the GSV at long-term follow-up. CAC was also associated with sustained improvements in symptoms and QoL, lower CEAP class, and high level of patient satisfaction without serious adverse effects between 36 and 60 months. The limitations of this publication included the small rate of successful follow up i.e., 36% of the original study randomized population, which could have introduced biases in the findings.

A systematic review by Dimech and Cassar (2020) was performed to assess the efficacy of n-butyl-2-cyanoacrylate (NBCA) glue in ablating primary truncal varicose veins and eliminating reflux compared with existing endovascular techniques. Secondary outcomes include complications and quality of life. PRISMA was used as a guide, and studies were screened for risk of bias and methodological quality. Subjects had to be  $\geq 18$  years of age and followed-up post-treatment with color Duplex ultrasound (DUS). Eligibility criteria included saphenofemoral junction (SFJ) or saphenopopliteal junction (SPJ) incompetence with reflux down truncal veins lasting  $> 0.5$  seconds on DUS interrogation and a Clinical, Etiological, Anatomical, and Pathophysiological classification of venous disorders ranging between C1 and C6. Out of 2,910 patients (3,220 veins) in 17 studies, 1,981 were administered NBCA, 445 radiofrequency ablation (RFA), and 484 EVLA with mean procedure times of 25.7, 23.2, and 28.7 minutes, respectively. Mean recruitment period was nine months (1-36 months) and followed-up for an average of 12.3 months (1-36 months). The majority were C2 to C3. Two-year occlusion rates were 93.7, 90.9, and 91.5% for NBCA, RFA, and EVLA, respectively. NBCA-treated patients experienced the least complications, with bruising, phlebitis, and pain being the most prevalent. Quality of life improved equally in all three modalities. The authors concluded that NBCA is simple to administer, safe, and effective even without compression stockings. The review was limited by lack of randomization for most included studies, and inclusion of products not currently FDA-approved. Further studies are required to assess longer-term benefit and the effect of anticoagulation on vein obliteration.

Gibson et al. (2018b) reported three-month outcomes from a post-market case series study of endovenous cyanoacrylate closure by the VenaSeal system (the WAVES study). Fifty subjects with symptomatic GSV, SSV, and/or accessory saphenous vein incompetence were treated with the VenaSeal system with no post procedure compression stockings. Concomitant procedures were not allowed as part of the original study protocol. Treating physicians predicted the type and nature of any concomitant procedures that they would usually perform at the time of ablation, if not limited by the constraints of the study. Evaluations were performed at one week, one and three months and included duplex ultrasound, numeric pain rating scale, revised VCSS, the AVVQ, and time to return to work and normal activities. At the three-month visit, the need for and type of adjunctive procedures were recorded. Complete closure at three months was achieved in 70 (99%) of the treated veins (48 GSVs, 14 accessory saphenous veins, eight SSVs). Revised VCSS improved from  $6.4 \pm 2.2$  to  $1.8 \pm 1.5$  ( $p < .001$ ) and AVVQ from  $17.3 \pm 7.9$  to  $6.5 \pm 7.2$  ( $p < .0001$ ). Sixty-six percent of patients underwent tributary treatment at three months. The percentage of patients who required adjunctive treatments at three months was lower than had been predicted by the treating physicians (65% versus 96%,  $p = .0002$ ). The authors reported that closure rates were high in the absence of the use of compression stockings or side branch treatment. Improvement in QoL was significant, and the need for and extent of concomitant procedures was significantly less than had been predicted by the treating physicians. Additional studies with larger patient

populations are needed to further evaluate the need for concomitant procedures with the VenaSeal system. These findings are limited by lack of comparison group undergoing a different treatment. This study was also included in the Hayes report (2022).

Gibson and Ferris (2017b) reported results of a prospective case series study (the WAVES study) of cyanoacrylate closure for the treatment of GSVs, SSVs, and/or accessory saphenous veins up to 20 mm in diameter (n = 50). Compression stockings post-procedure were not utilized. Patients returned at one week and one month for follow-up. All treated veins (48 GSVs, 14 accessory saphenous veins, and eight SSVs) had complete closure by duplex ultrasound at seven days and one month. Mean time to return to work and normal activities was 0.2 ±1.1 and 2.4 ±4.1 days, respectively. The revised VCSS was improved to 1.8 ±1.4 (p < .001) and AVVQ score to 8.9 ±6.6 (p < .001) at one month. Phlebitis in the treatment area or side branches occurred in 10 subjects (20%) and completely resolved in all but one subject (2%) by one month. The authors concluded that cyanoacrylate closure is safe and effective for the treatment of one or more incompetent saphenous or accessory saphenous veins, closure rates were high even in the absence of the use of compression stockings or side branch treatment. Time back to work or normal activities was short and improvements in venous severity scores and QoL were in the authors' opinion significant, comparing favorably with alternative treatment methods. RCTs with a larger patient population and longer follow-up periods are needed to validate findings. The findings of this study are limited by lack of comparison group undergoing a different treatment approach. This study was also included in the Hayes report (2022).

Almeida et al. (2015) evaluated the safety and effectiveness of endovenous cyanoacrylate-based embolization of incompetent GSVs in a case series study of 38 patients. At 12 months, 36 patients were available for follow-up and 24 patients at 24 months. Complete occlusion of the treated GSV was confirmed by duplex ultrasound in all patients except for one complete and two partial recanalizations observed at one, three and six months of follow-up, respectively. Kaplan-Meier analysis yielded an occlusion rate of 92.0% (95% CI 0.836-1.0) at 24 months follow-up. VCSS improved in all patients from a mean of 6.1 ±2.7 at baseline to 1.3 ±1.1, 1.5 ±1.4 and 2.7 ±2.5 at six, 12 and 24 months, respectively (p < .0001). Edema improved in 89% of legs (n = 34) at 48 hours follow-up. At baseline, only 13% were free from pain. At six, 12 and 24 months, 84%, 78% and 64% were free from leg pain, respectively. In a follow-up study, Almeida et al. (2017) evaluated the long-term safety and effectiveness of endovenous cyanoacrylate (CA)-based closure of incompetent GSV on the twenty-nine individuals that were available for the 36-month follow-up. Complete occlusion of the treated veins was confirmed by ultrasound in all subjects with the exception of two subjects showing recanalization at month one and month three. Kaplan-Meier analysis revealed an occlusion rate at month 36 of 94.7%. The mean VCSS improved from 6.1 ± 2.7 at baseline to 2.2 ± 0.4 at month 36 (p < .0001). Pain, edema, and varicosities (VCSS subdomains) improved in 75.9%, 62.1%, and 41.4% of subjects, respectively, at month 36. Overall adverse events were self-limited and mild or moderate. The authors concluded cyanoacrylate adhesive had no reported serious adverse events, had long-term occlusion rates comparable to other thermal and nonthermal methods, and appears to be safe and effective for saphenous vein closure. Small sample size and lack of comparison groups are limitations to this study.

A prospective multicenter case series study was conducted on 78 patients with GSV reflux using cyanoacrylate embolization (Proebstle et al., 2015). Clinical examination, QoL assessment and duplex ultrasound were performed at two days, one, three, six, and 12 months. 68 (97.1%) were available for 12-month follow-up. Two-day follow-up showed one proximal and one distal partial recanalization. Three additional proximal recanalizations were observed at 3-month (n = 2) and 6-month (n = 1) follow-up. Cumulative 12-month survival free from recanalization was 92.9% (95% confidence interval, 87.0%-99.1%). Mean (standard deviation) VCSS improved from 4.3 ±2.3 at baseline to 1.1 ±1.3 at 12 months. AVVQ score showed an improvement from 16.3 at baseline to 6.7 at 12 months (p < .0001). Side effects were generally mild; a phlebotic reaction occurred in eight cases (11.4%) with a median duration of 6.5 days (range, 2-12 days). Pain without a phlebotic reaction was observed in five patients (8.6%) for a median duration of 1 day (range, 0-12 days). No serious adverse event occurred. Paresthesia was not observed. The authors concluded that endovenous CA embolization of refluxing GSVs is safe and effective without the use of tumescent anesthesia or compression stockings. Additional studies are needed to validate the effectiveness of cyanoacrylate embolization.

## VenoValve

Evidence in peer review literature evaluating VenoValve porcine bioprosthetic valve for the treatment of chronic venous insufficiency is limited. Future robust RCTs are warranted along with long-term outcomes to establish the safety and efficacy of this procedure.

A 2022 Hayes Emerging Technology Report states published evidence is limited to publications reporting 6-month and 1-year outcomes for 11 patients. The VenoValve will be the first porcine bioprosthetic valve to reach the market in the U.S., and the

first device approved to treat CVI, if eventually FDA-approved. VenoValve is currently under investigation in the Surgical Anti-Reflux Venous Valve Endoprosthesis (SAVVE) trial (NCT04943172).

Ulloa and Glickman (2021) conducted a single-center, prospective, non-randomized, first-in-human trial using a prosthetic venous valve, VenoValve, for patients with severe chronic venous insufficiency (C4b-C6 disease). Ten patients had the prosthetic valve surgically implanted into the femoral vein. Follow-up examinations were conducted postoperatively at two and 14 days and then every 30 days for six months to evaluate feasibility, initial safety, and performance outcomes of the VenoValve. Six patients had required bovine patch angioplasty of the vein. Four adverse events occurred, including one case of hematoma at the incision site that was aspirated, two cases of superficial wound infection in C6 patients treated with antibiotics, and one case of a bleeding complication due to warfarin anticoagulation. One patient's VenoValve had thrombosed at five months due to nontherapeutic anticoagulation. Improvements in all five patients who had reached the 6-month follow-up mark with the VenoValve were demonstrated during the study period by decreases in the VCSS (61% decrease from baseline), visual analog scale for pain scores (57% decrease), and reflux time (40% decrease) and a statistically significant improvement in the VEINES-QOL/Sym questionnaire. The patient with the occluded VenoValve had experienced improvements in all areas except for the reflux time. The authors concluded that VenoValve showed promising results with improvements noted in QOL and clinical outcomes. The authors recommended further follow-up and larger studies in the future.

## **Clinical Practice Guidelines**

### ***American College of Phlebology***

The American College of Phlebology Guidelines Committee (Gibson et al., 2017c) performed a systematic review of the literature regarding the clinical impact and treatment of incompetent accessory saphenous veins. They developed a consensus opinion that patients with symptomatic incompetence of the accessory great saphenous veins (anterior and posterior accessory saphenous veins) be treated with EVTA (laser or radiofrequency) or UGFS to eliminate symptomatology (Recommendation Grade 1C).

The American College of Phlebology Guidelines Committee (2016) updated their evidence-based recommendations for treatment of superficial venous disease of the lower leg. They recommend that named veins (GSV, SSV, AAGSV, posterior accessory of the great saphenous vein [PAGSV], intersaphenous vein [Vein of Giacomini]) must have a reflux time > 500 msec regardless of the reported vein diameter (Grade 1A).

EVTA (laser and radiofrequency) is the Committee's preferred treatment for saphenous and accessory saphenous (GSV, SSV, AAGSV, PAGSV) vein incompetence (Grade 1B). They suggest mechanical/chemical ablation may also be used to treat truncal venous reflux (Grade 2B). They further comment that open surgery is appropriate in veins not amenable to endovenous procedures but otherwise is not recommended because of increased pain, convalescent time, and morbidity (Grade 1B).

### ***European Society for Vascular Surgery (ESVS)***

The ESVS released a clinical practice guideline for management of chronic venous disease (De Maeseneer et al., 2022). The guidelines state that for patients with GSV and SSV incompetence requiring treatment endovenous thermal ablation is recommended as the first-choice treatment, in preference to high ligation/stripping and USGF. However, USGF may be considered for treating saphenous trunks with a diameter less than 6mm. The guidelines note that in long term follow up of comparative studies, treatment with UGFS has been substantially less effective than EVLA, RFA, and surgery in terms of occlusion or absence rates. Additionally, foam sclerotherapy is the technique of choice for anatomical configurations that make endovenous cannulation or advancing the ablation device challenging, and is suitable for treating tortuous, recurrent varicose veins. Mechanochemical ablation and cyanoacrylate adhesive closure may be considered when a non-thermal technique is preferred for patients with GSV incompetence. For patients with GSV incompetence, high ligation/stripping should be considered, if endovenous thermal ablation options are not available. Endovenous non-thermal non tumescent ablation methods may be considered for treatment of SSV incompetence. Additionally, endovenous thermal ablation and USGF may be considered for anterior accessory saphenous vein requiring treatment.

### ***National Institute for Health and Care Excellence (NICE)***

In 2020, the National Institute for Health and Care Excellence (NICE) released an update to their guidance on Cyanoacrylate Glue Occlusion for Varicose Veins. The updated guidance states that current evidence on the safety and efficacy of cyanoacrylate glue occlusion for varicose veins is adequate to support the use of this procedure provided that standard

arrangements are in place for clinical governance, consent, and audit. In addition, the guideline states physicians should: 1) only perform the procedure after appropriate training and experience in the use of venous ultrasound; 2) discuss the available options with the patient before making a decision; and 3) follow their hospital's policies regarding performing procedures and monitoring results.

In an updated guideline on endovenous MOCA for varicose veins, NICE (2016) states that current evidence on the safety and efficacy of endovenous MOCA for varicose veins appears adequate to support the use of this procedure provided that standard arrangements are in place for consent, audit, and clinical governance. Clinicians are encouraged to collect longer-term follow-up data.

The NICE 2013 interventional procedure guidance on UGFS specifies that if symptoms related to varicose veins are severe, the main treatment options include endovenous laser treatment and radiofrequency ablation, and surgery (ligation and stripping of the GSVs or ligation with or without stripping of the SSVs, and phlebectomy). The NICE 2013 clinical guideline on the diagnosis and treatment of varicose veins adds that if endovenous ablation is unsuitable, offer UGFS.

### ***Society for Vascular Surgery (SVS)/American Venous Forum (AVF)/American Vein and Lymphatic Society (AVLS)/Society of Interventional Radiology (SIR)***

The SVS, AVF, and AVLS collaborated to update the 2011 SVS/AVF clinical practice guideline to provide evidence-based recommendations for treating patients with varicose veins of the lower limbs (Gloviczki. et al., 2022). Recommendations of the guideline are summarized as follows (not all-inclusive):

- For patients with CVD of the lower extremities, duplex ultrasound scanning is the diagnostic test of choice for evaluation of venous reflux
- Reflux is defined as a minimum value > 500 ms of reversed flow in the superficial truncal veins and the tibial, deep femoral, and perforating veins
- Axial reflux is defined as uninterrupted retrograde venous flow from the groin to the calf, and junctional reflux is limited to the SFJ or SPJ
- Use of the 2020 upgraded CEAP classification of chronic venous disorders is recommended
- “Pathologic” perforating veins in patients with varicose veins (CEAP clinical class C2) includes those with an outward flow duration of ≥ 500 ms and a diameter of ≥ 3.5 mm on duplex ultrasound
- For patients with symptomatic varicose veins and axial reflux in the GSV and SSV, treatment with endovenous ablation over high ligation and stripping is recommended due to less postprocedure pain and morbidity, and an earlier return to regular activity; if the technology or expertise in endovenous ablation is not available or the venous anatomy precludes endovenous treatment, ligation and stripping is recommended
- For patients with symptomatic varicose veins and axial reflux in the GSV, SSV, who place a high priority on the long-term outcomes of treatment (QoL and recurrence), treatment with EVLA, RFA, or high ligation and stripping over physician compounded UGFS is suggested
- For patients with symptomatic axial reflux, both thermal and nonthermal ablation of the GSV and SSV are recommended Depending on the available expertise of the treating physician and the preference of the patient
- In patients with symptomatic reflux in the GSV or SSV and associated varicosities, ablation of the refluxing venous trunk, and concomitant phlebectomy, or UGFS of the varicosities with physician-compounded foam or commercial PEM is recommended
- In patients with symptomatic reflux in the GSV or SSV, ablation of the refluxing venous trunk, and staged or UGFS of the varicosities is recommended only if anatomic or medical reasons are present

The SVS, AVF, AVLS, and SIR developed the appropriate use criteria (AUC) for chronic lower extremity venous disease using the RAND/UCLA Appropriateness Method incorporating best available evidence with expert opinion and engaging a panel of experts in the field through a modified Delphi exercise (Masuda et al. 2020). The consensus does not appear to be based on a systematic review of the literature. One hundred and nineteen scenarios were rated on a scale of one to nine by an expert panel, with one being never appropriate and nine being appropriate. The panelists rated ablation for axial reflux of the GSV, with or without SFJ reflux, in symptomatic patients, CEAP classes 2-6 as appropriate. Per the AUC, when accompanied by no SFJ reflux (the junction is either assumed or proven to be competent or previously interrupted and communicates with the GSV through incompetent thigh perforators or other sources of collateral flow) the remaining refluxing GSV may be the source of recurrent symptoms. Therefore, for axial GSV reflux, ablating the GSV will likely lead to decreased recurrence even if the SFJ

shows no reflux. The authors note that the AUC statements were intended to serve as a guide to patient care, particularly in areas where high quality evidence is lacking and was not meant to be a guide that addresses all clinical situations.

The SVS and AVF released joint clinical practice guidelines regarding the care of patients with venous leg ulcers (O'Donnell et al., 2014). For patients with a venous leg ulcer (C6), and incompetent superficial veins that have reflux to the ulcer bed in addition to pathological perforating veins (> 500 ms reflux duration and diameter of > 3.5 mm), that are located beneath or associated with the ulcer bed, the guideline recommends ablation of both the incompetent superficial veins and perforator veins in addition to standard compressive therapy to aid in ulcer healing and prevent recurrence. For patients who are at risk for a venous leg ulcer (C4b), or have a healed venous ulcer (C5), and have axial reflux directed to the bed of the affected skin/ulcer, the guidelines recommend ablation of the incompetent superficial veins in addition to standard compressive therapy.

## U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Vein ligation surgery is a procedure and therefore not subject to FDA regulation.

The ClariVein® infusion catheter (Vascular Insights) received FDA approval (K071468) on March 20, 2008. The device is designed to introduce physician-specified medicaments into the peripheral vasculature. Refer to the following website for more information: [http://www.accessdata.fda.gov/cdrh\\_docs/pdf7/K071468.pdf](http://www.accessdata.fda.gov/cdrh_docs/pdf7/K071468.pdf). (Accessed June 6, 2023)

The VenaSeal™ Closure System received the FDA's pre-market approval (PMA) on February 20, 2015 (P140018). The device is indicated for the permanent closure of lower extremity superficial truncal veins, such as the great saphenous vein (GSV), through endovascular embolization with coaptation. VenaSeal is intended for use in adults with clinically symptomatic venous reflux as diagnosed by duplex ultrasound (DUS). Refer to the following website for more information: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P140018>. (Accessed June 6, 2023)

Varithena (polidocanol injectable foam) (Provensis Ltd.) received FDA approval on November 25, 2013, as a sclerosing agent indicated for the treatment of incompetent great saphenous veins, accessory saphenous veins and visible varicosities of the great saphenous vein system above and below the knee. Refer to the following websites for more information:

- [https://www.accessdata.fda.gov/drugsatfda\\_docs/applletter/2013/205098Orig1s000ltr.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/applletter/2013/205098Orig1s000ltr.pdf)
- [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2013/205098s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2013/205098s000lbl.pdf)

(Accessed June 6, 2023)

## References

Almeida JI, Javier JJ, Mackay EG, et al. Thirty-sixth-month follow-up of first-in-human use of cyanoacrylate adhesive for treatment of saphenous vein incompetence. *J Vasc Surg Venous Lymphat Disord*. 2017 Sep;5(5):658-666.

Almeida JI, Javier JJ, Mackay EG, et al. Two-year follow-up of first human use of cyanoacrylate adhesive for treatment of saphenous vein incompetence. *Phlebology*. 2015 Jul;30(6):397-404.

American College of Phlebology. Practice guidelines. Superficial venous disease. Treatment of superficial venous disease of the lower leg. October 2014. Updated February 2016.

American Medical Association. Current Procedural Terminology: CPT Professional Edition. AMA Press.

Amshar M, Nugraha RA, Batubara EAD, et al. Cyanoacrylate embolization versus endovenous laser ablation in treating saphenous vein insufficiency: a systematic review and meta-analysis. *Ann Vasc Surg*. 2022 Mar;80:313-324.

Bishawi M, Bernstein R, Boter M, et al. Mechanochemical ablation in patients with chronic venous disease: A prospective multicenter report. *Phlebology*. 2014 Jul;29(6):397-400.

Boersma D, Kornmann VN, van Eekeren RR, et al. Treatment modalities for small saphenous vein insufficiency: systematic review and meta-analysis. *J Endovasc Ther*. 2016 Feb;23(1):199-211.

Boersma D, van Eekeren RR, Werson DA, et al. Mechanochemical endovenous ablation of small saphenous vein insufficiency using the ClariVein® device: one-year results of a prospective series. *Eur J Vasc Endovasc Surg.* 2013 Mar;45(3):299-303.

Bootun R, Lane T, Dharmarajah B, et al. Intra-procedural pain score in a randomised controlled trial comparing mechanochemical ablation to radiofrequency ablation: The multicentre Venefit™ versus ClariVein® for varicose veins trial. *Phlebology.* 2016 Feb;31(1):61-5.

Bozkurt AK, Yilmaz MF. A prospective comparison of a new cyanoacrylate glue and laser ablation for the treatment of venous insufficiency. *Phlebology.* 2016 Mar;31(1 Suppl):106-13.

ClariVein website. Vascular Insights, LLC. Available at: <http://clarivein.com/>. Accessed June 6, 2023.

CPT Assistant. Correct coding of endovascular treatment for low extremity venous incompetency. October 2014.

CPT Assistant. Therapy for incompetent veins. November 2016.

De Maeseneer MG, Kakkos SK, Aherne T, et al. Editor's Choice - European Society for Vascular Surgery (ESVS) 2022 Clinical Practice Guidelines on the management of chronic venous disease of the lower limbs. *Eur J Vasc Endovasc Surg.* 2022 Feb;63(2):184-267.

Dimech AP, Cassar K. Efficacy of cyanoacrylate glue ablation of primary truncal varicose veins compared to existing endovenous techniques: a systematic review of the literature. *Surg J (N Y).* 2020 Jun 19;6(2):e77-e86.

ECRI Institute. Varithena Injectable Foam (Boston Scientific Corp.) for treating varicose veins. Plymouth Meeting (PA): ECRI; 2020 Dec 11. (Clinical Evidence Assessment).

ECRI. VenaSeal Closure System (Medtronic plc.) for embolizing varicose veins. Plymouth Meeting (PA): ECRI; 2021 Feb 4. (Clinical Evidence Assessment).

Elias S, Raines JK. Mechanochemical tumescentless endovenous ablation: final results of the initial clinical trial. *Phlebology.* 2012 Mar;27(2):67-72.

Eroglu E, Yasim A. A randomised clinical trial comparing n-butyl cyanoacrylate, radiofrequency ablation and endovenous laser ablation for the treatment of superficial venous incompetence: two year follow up results. *Eur J Vasc Endovasc Surg.* 2018 Oct;56(4):553-560.

Flessenkämper I, Hartmann M, Hartmann K et al. Endovenous laser ablation with and without high ligation compared to high ligation and stripping for treatment of great saphenous varicose veins: Results of a multicentre randomized controlled trial with up to 6 years follow-up. *Phlebology.* 2016 Feb;31(1):23-33.

Gibson K, Ferris B. Cyanoacrylate closure of incompetent great, small and accessory saphenous veins without the use of post-procedure compression: Initial outcomes of a post-market evaluation of the VenaSeal System (the WAVES Study). *Vascular.* 2017b Apr;25(2):149-156.

Gibson K, Kabnick L; Varithena® 013 Investigator Group. A multicenter, randomized, placebo-controlled study to evaluate the efficacy and safety of Varithena® (polidocanol endovenous microfoam 1%) for symptomatic, visible varicose veins with saphenofemoral junction incompetence. *Phlebology.* 2017a Apr;32(3):185-193.

Gibson K, Khilnani N, Schul M, et al. American College of Phlebology Guidelines - treatment of refluxing accessory saphenous veins. *Phlebology.* 2017c Aug;32(7):448-452.

Gibson K, Minjarez R, Gunderson K, et al. Need for adjunctive procedures following cyanoacrylate closure of incompetent great, small and accessory saphenous veins without the use of postprocedure compression: Three-month data from a postmarket evaluation of the VenaSeal System (the WAVES Study). *Phlebology.* 2018b Sep 18:268355518801641.

Gibson K, Morrison N, Kolluri R, et al. Twenty-four-month results from a randomized trial of cyanoacrylate closure versus radiofrequency ablation for the treatment of incompetent great saphenous veins. *J Vasc Surg Venous Lymphat Disord.* 2018a Sep;6(5):606-613.

Gloviczki P, Lawrence PF, Wasan SM, et al. The 2022 Society for Vascular Surgery, American Venous Forum, and American Vein and Lymphatic Society clinical practice guidelines for the management of varicose veins of the lower extremities. Part I. Duplex scanning and treatment of superficial truncal reflux: Endorsed by the Society for Vascular Medicine and the International Union of Phlebology. *J Vasc Surg Venous Lymphat Disord.* 2022 Oct 12:S2213-333X(22)00417-6.

Hayes, Inc. Hayes Emerging Technology Report. VenoValve for chronic venous insufficiency. Lansdale, PA: Hayes, Inc.; May 2022.

Hayes, Inc. Hayes Health Technology Assessment. Cyanoacrylate Embolization With the VenaSeal Closure System (Medtronic Inc.) for the Treatment of Varicose Veins Cyanoacrylate embolization (VenaSeal Closure System) for the treatment of varicose veins.: Hayes, Inc.; December 2020. Updated December 2021.

Hayes, Inc. Hayes Health Technology Assessment. Mechanochemical endovenous ablation (MOCA) with the ClariVein Infusion Catheter Nonthermal Vein Ablation System (Merit Medical Systems Inc.) for treatment of varicose veins.: Hayes, Inc.; February 2022.

Hayes, Inc. Hayes Health Technology Brief. Polidocanol endovenous microfoam (Varithena) 1% for treatment of varicose veins. Hayes, Inc.; May 2016. Updated November 2022.

Holewijn S, van Eekeren RRJP, Vahl A, et al. Two-year results of a multicenter randomized controlled trial comparing mechanochemical endovenous ablation to radiofrequency ablation in the treatment of primary great saphenous vein incompetence (MARADONA trial). *J Vasc Surg Venous Lymphat Disord*. 2019 May;7(3):364-374.

Joh JH, Lee T, Byun SJ, et al. A multicenter randomized controlled trial of cyanoacrylate closure and surgical stripping for incompetent great saphenous veins. *J Vasc Surg Venous Lymphat Disord*. 2022 Mar;10(2):353-359.

Jones L, Braithwaite BD, Selwyn D, et al. Neovascularisation is the principal cause of varicose vein recurrence: results of a randomized trial of stripping the long saphenous vein. *Eur J Vasc Endovasc Surg*. 1996 Nov;12(4):442-5.

Kim PS, Bishawi M, Draughn D, et al. Mechanochemical ablation for symptomatic great saphenous vein reflux: a two-year follow-up. *Phlebology*. 2017;32(1):43-48.

King JT, O'Byrne M, Vasquez M, et al. Treatment of truncal incompetence and varicose veins with a single administration of a new polidocanol endovenous microfoam preparation improves symptoms and appearance. *Eur J Vasc Endovasc Surg*. 2015 Dec;50(6):784-93.

Kolluri R, Gibson K, Cher D, et al. Roll-in phase analysis of clinical study of cyanoacrylate closure for incompetent great saphenous veins. *J Vasc Surg Venous Lymphat Disord*. 2016 Oct;4(4):407-15.

Lane T, Bootun R, Dharmarajah B, et al. A multi-centre randomised controlled trial comparing radiofrequency and mechanical occlusion chemically assisted ablation of varicose veins – Final results of the Venefit versus Clarivein for varicose veins trial. *Phlebology*. 2017 Mar;32(2):89-98.

Lawaetz M, Serup J, Lawaetz B, et al. Comparison of endovenous ablation techniques, foam sclerotherapy and surgical stripping for great saphenous varicose veins. Extended 5-year follow-up of a RCT. *Int Angiol*. 2017 Jun;36(3):281-288.

Lurie F, Passman M, Meisner M, et al. The 2020 update of the CEAP classification system and reporting standards. *J Vasc Surg Venous Lymphat Disord*. 2020 May;8(3):342-352. Erratum in: *J Vasc Surg Venous Lymphat Disord*. 2021 Jan;9(1):288.

Marston WA, Owens LV, Davies S, et al. Endovenous saphenous ablation corrects the hemodynamic abnormality in patients with CEAP clinical class 3-6 CVI due to superficial reflux. *Vasc Endovascular Surg*. 2006 Mar-Apr;40(2):125-30.

Masuda E, Ozsvath K, Vossler J, et al. The 2020 appropriate use criteria for chronic lower extremity venous disease of the American Venous Forum, the Society for Vascular Surgery, the American Vein and Lymphatic Society, and the Society of Interventional Radiology. *J Vasc Surg Venous Lymphat Disord*. 2020 Jul;8(4):505-525.e4.

Mohamed AH, Leung C, Wallace T, et al. A randomized controlled trial of endovenous laser ablation versus mechanochemical ablation with Clarivein in the management of superficial venous incompetence (LAMA trial). *Ann Surg*. 2021 Jun 1;273(6):e188-e195.

Morrison N, Gibson K, McEnroe S, et al. Randomized trial comparing cyanoacrylate embolization and radiofrequency ablation for incompetent great saphenous veins (VeClose). *J Vasc Surg*. 2015 Apr;61(4):985-94.

Morrison N, Gibson K, Vasquez M, et al. Five-year extension study of patients from a randomized clinical trial (VeClose) comparing cyanoacrylate closure versus radiofrequency ablation for the treatment of incompetent great saphenous veins. *J Vasc Surg Venous Lymphat Disord*. 2020 Nov;8(6):978-989.

Morrison N, Gibson K, Vasquez M, et al. VeClose trial 12-month outcomes of cyanoacrylate closure versus radiofrequency ablation for incompetent great saphenous veins. *J Vasc Surg Venous Lymphat Disord*. 2017 May;5(3):321-330.

Morrison N, Kolluri R, Vasquez M, et al. Comparison of cyanoacrylate closure and radiofrequency ablation for the treatment of incompetent great saphenous veins: 36-month outcomes of the VeClose randomized controlled trial. *Phlebology*. 2019 Jul;34(6):380-390.2018.



National Heart, Lung and Blood Institute (NHLBI). Varicose veins. February 2014. Available at: <https://www.nhlbi.nih.gov/health-topics/varicose-veins>. Accessed June 6, 2023.

National Institute for Health and Care Excellence (NICE). Clinical guideline (CG168). Varicose veins: diagnosis and management. July 2013.

National Institute for Health and Care Excellence (NICE). Cyanoacrylate glue occlusion for varicose veins. Interventional procedure guidance (IPG670). March 2020.

National Institute for Health and Care Excellence (NICE). Interventional procedures guidance (IPG557). Endovenous mechanochemical ablation for varicose veins. May 2016.

National Institute for Health and Care Excellence (NICE). Interventional procedures guidance (IPG 440). Ultrasound-guided foam sclerotherapy for varicose veins. February 2013.

O'Donnell TF Jr, Passman MA, Marston WA, et al.; Society for Vascular Surgery; American Venous Forum. Management of venous leg ulcers: clinical practice guidelines of the Society for Vascular Surgery® and the American Venous Forum. *J Vasc Surg*. 2014 Aug;60(2 Suppl):3S-59S.

Proebstle TM, Alm J, Dimitri S, et al. The European multicenter cohort study on cyanoacrylate embolization of refluxing great saphenous veins. *J Vasc Surg Venous Lymphat Disord*. 2015 Jan;3(1):2-7.

Ulloa JH, Glickman M. Human trial using the novel bioprosthetic VenoValve in patients with chronic venous insufficiency. *J Vasc Surg Venous Lymphat Disord*. 2021 Jul;9(4):938-944.

Vähäaho S, Mahmoud O, Halmesmäki K, et al. Randomized clinical trial of mechanochemical and endovenous thermal ablation of great saphenous varicose veins. *Br J Surg*. 2019 Apr;106(5):548-554.

van Eekeren RR, Boersma D, Elias S, et al. Endovenous mechanochemical ablation of great saphenous vein incompetence using the ClariVein device: a safety study. *J Endovasc Ther*. 2011 Jun;18(3):328-34.

van Eekeren RR, Boersma D, Konijn V, et al. Postoperative pain and early quality of life after radiofrequency ablation and mechanochemical endovenous ablation of incompetent great saphenous veins. *J Vasc Surg*. 2013 Feb;57(2):445-50.

Varithena® website. Boston Scientific. Available at: <https://www.varithena.com/en-us/home.html>. Accessed June 6, 2023.

VenaSeal™ Closure System website. Medtronic. Available at: <https://www.medtronic.com/us-en/patients/treatments-therapies/varicose-vein-therapies/our-treatments/venaseal-closure-system.html>. Accessed June 6, 2023.

Vos CG, Ünlü C, Bosma J, et al. A systematic review and meta-analysis of two novel techniques of nonthermal endovenous ablation of the great saphenous vein. *J Vasc Surg Venous Lymphat Disord*. 2017 Nov;5(6):880-896.

Vun SV, Rashid ST, Blest NC, Spark JL. Lower pain and faster treatment with mechanico-chemical endovenous ablation using ClariVein®. *Phlebology*. 2015 Dec;30(10):688-92.

Winterborn, RJ, Foy C, Earnshaw JJ. Causes of varicose vein recurrence: late results of a randomized controlled trial of stripping the long saphenous vein. *J Vasc Surgery*. 2004 Oct 40(4): 634-9.

Witte ME, Holewijn S, van Eekeren RR, et al. Midterm outcome of mechanochemical endovenous ablation for the treatment of great saphenous vein insufficiency. *J Endovasc Ther*. 2017b Feb;24(1):149-155.

Witte ME, Zeebregts CJ, de Borst GJ, et al. Mechanochemical endovenous ablation of saphenous veins using the ClariVein: A systematic review. *Phlebology*. 2017a Dec;32(10):649-657.

Wittens C, Davies AH, Bækgaard N, et al. Editor's choice - management of chronic venous disease: clinical practice guidelines of the European Society for Vascular Surgery (ESVS). *Eur J Vasc Endovasc Surg*. 2015 Jun;49(6):678-737.

## Policy History/Revision Information

Date	Summary of Changes
12/01/2023	<p><b>Definitions</b></p> <ul style="list-style-type: none"> <li>● Removed definition of:               <ul style="list-style-type: none"> <li>○ Cosmetic Procedures</li> <li>○ Reconstructive Surgery</li> </ul> </li> </ul>

Date	Summary of Changes
	<p><b>Applicable Codes</b></p> <ul style="list-style-type: none"> <li>Removed CPT codes 37760 and 37761</li> </ul> <p><b>Supporting Information</b></p> <ul style="list-style-type: none"> <li>Updated <i>References</i> section to reflect the most current information</li> <li>Archived previous policy version CS117IN.07</li> </ul>

## Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual<sup>®</sup> criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.