

Preventive Care Services

Policy Number: RMHP.MP.016.52
Effective Date: March 1, 2025

[➔ Instructions for Use](#)

Table of Contents	Page
Coverage Rationale	1
Frequently Asked Questions	3
Definitions	5
Applicable Codes	5
Benefit Considerations	45
Clinical Evidence	45
References	45
Instructions for Use	45

Related Policies
• Breast Imaging for Screening and Diagnosing Cancer
• Cardiovascular Disease Risk Tests
• Colonoscopy Guidelines
• Consultation Services
• Cytological Examination of Breast Fluids for Cancer Screening or Diagnosis
• Genetic Testing for Hereditary Cancer
• Long-Acting Injectable Antiretroviral Agents for HIV
• Magnetic Resonance Imaging (MRI) and Computed Tomography (CT) Scan – Site of Service
• Outpatient Surgical Procedures – Site of Service
• Preventive Medicine and Screening
• Preventive Vaccines (Immunizations)
• Screening Colonoscopy Procedures – Site of Service

Coverage Rationale

Indications for Coverage

Introduction

UnitedHealthcare covers certain medical services under the preventive care services benefit. The federal Patient Protection and Affordable Care Act (PPACA) requires non-grandfathered health plans to cover certain “recommended preventive services” as identified by PPACA under the preventive care services benefit, without cost sharing to members when provided by network providers. This includes:

- Evidence-based items or services that have in effect a rating of “A” or “B” in the current recommendations of the United States Preventive Services Task Force.
- Immunizations for routine use in children, adolescents and adults that have in effect a recommendation from the *Advisory Committee on Immunization Practices of the Centers for Disease Control and Prevention*.
- With respect to infants, children and adolescents, evidence-informed preventive care and screenings provided for in the comprehensive guidelines supported by the *Health Resources and Services Administration*.
- With respect to women, such additional preventive care and screenings as provided for in comprehensive guidelines supported by the *Health Resources and Services Administration*.

Member Cost-Sharing

Non-Grandfathered Plans

- Non-grandfathered plans provide coverage for preventive care services with no member cost sharing (i.e., covered at 100% of Allowed Amounts without deductible, coinsurance, or copayment) when services are obtained from a Network provider.
- Under PPACA, services obtained from an out-of-network provider are not required to be covered under a plan’s preventive benefit and may be subject to member cost sharing. Refer to the member specific benefit plan document for out-of-network benefit information, if any.

Grandfathered Plans

- Plans that maintain grandfathered status under PPACA are not required by law to provide coverage for these preventive services without member cost sharing; although a grandfathered plan may choose to voluntarily amend its plan document to include these preventive benefits.
- Except where there are state mandates, a grandfathered plan might include member cost sharing, or exclude some of the preventive care services identified under PPACA.
- Refer to the member specific benefit plan document for details on how benefits are covered under a grandfathered plan.

Preventive vs. Diagnostic Services

Certain services can be done for preventive or diagnostic reasons. When a service is performed for the purpose of preventive screening and is appropriately reported, it will be considered under the preventive care services benefit. This includes services directly related to the performance of a covered preventive care service (refer to the [Frequently Asked Questions](#) section for additional information.)

Preventive services are those performed on a person who:

- has not had the preventive screening done before and does not have symptoms or other abnormal studies suggesting abnormalities*; or
- has had screening done within the recommended interval with the findings considered normal; or
- has had diagnostic services results that were normal after which the physician recommendation would be for future preventive screening studies using the preventive services intervals.

*In the case of a colonoscopy done as a follow-up to a positive stool-based screening (e.g., FIT, FOBT, and fecal DNA) or direct visualization screening (e.g., sigmoidoscopy or CT colonography), refer to FAQ#4 below.

When a service is done for diagnostic purposes, it will be considered under the applicable non-preventive medical benefit. Diagnostic services are done on a person who:

- had abnormalities found on previous preventive or diagnostic studies that require further diagnostic studies; or
- had abnormalities found on previous preventive or diagnostic studies that would recommend a repeat of the same studies within shortened time intervals from the recommended preventive screening time intervals; or
- had a symptom(s) that required further diagnosis; or
- does not fall within the applicable population for a recommendation or guideline.

Covered Breastfeeding Equipment

Personal-use electric breast pump:

- The purchase of a personal-use electric breast pump (HCPCS code E0603).
 - This benefit is limited to one pump per birth. In the case of a birth resulting in multiple infants, only one breast pump is covered.
 - A breast pump purchase includes the necessary supplies for the pump to operate.
- Replacement breast pump supplies necessary for the personal-use electric breast pump to operate. This includes: standard power adaptor, tubing adaptors, tubing, locking rings, bottles specific to breast pump operation, caps for bottles that are specific to the breast pump, valves, filters, and breast shield and/or splash protector for use with the breast pump.
- Breastmilk storage bags (HCPCS code A4287).

Colonoscopies

Colonoscopy – Preventive Care Services Benefit (Without Member Cost-Sharing)

Member cost-sharing for colonoscopy is waived when all of the following apply:

- The patient's age is 45-75 years (ends on 76th birthday) as recommended by the USPSTF; and
- The provider is participating in the network; and
- When billed in accordance with the coding in the [Colorectal Cancer Screening](#) row listed in this policy.

Colonoscopy may require a site of service review. Refer to the Medical Policies titled [Screening Colonoscopy Procedures – Site of Service](#) and [Outpatient Surgical Procedures – Site of Service](#).

Colonoscopy – Medical Benefit (With Member Cost-Sharing)

Member cost-sharing may apply when a colonoscopy is done in any one of the following scenarios:

- The patient's age is outside of the age recommendation of the USPSTF (age 45-75 years); or

- The provider is non-network; or
- Colonoscopy performed with a shortened time interval outside of the USPSTF recommendations; or
- Colonoscopy performed for diagnostic purposes; or
- Colonoscopy performed for surveillance purposes (e.g., a follow-up colonoscopy performed after identification or removal of a polyp or cancer on a previous colonoscopy); or
- Colonoscopy performed for therapeutic/treatment purposes.

The above colonoscopies may require advanced notification and/or site of service review. Refer to the [Colonoscopy Guidelines](#) and the Medical Policy titled [Outpatient Surgical Procedures – Site of Service](#).

Coverage Limitations and Exclusions

- Services not covered under the preventive care benefit may be covered under another portion of the medical benefit plan.
- The coverage outlined in this policy does not address certain outpatient prescription medications, tobacco cessation drugs and/or over the counter items, as required by PPACA. These preventive benefits are administered by the member’s pharmacy plan administrator. For details on coverage, refer to the member-specific pharmacy plan administrator.
- A vaccine (immunization) is not covered if it does not meet company vaccine policy requirements for FDA labeling and if it does not have explicit ACIP recommendations for routine use published in the Morbidity and Mortality Weekly Report (MMWR) of the Centers for Disease Control and Prevention (CDC) and is not listed on the applicable immunization schedule of ACIP. (Refer to the [Preventive Care Services: Vaccine Codes](#).)
- Examinations, screenings, testing, or vaccines (immunizations) are not covered when:
 - required solely for the purposes of career or employment, school or education, sports or camp, travel [including travel vaccines (immunizations)], insurance, marriage, or adoption; or
 - related to judicial or administrative proceedings or orders; or
 - conducted for purposes of medical research; or
 - required to obtain or maintain a license of any type.
- Services that are investigational, experimental, unproven, or not medically necessary are not covered.
- Breastfeeding equipment and supplies not listed above. This includes, but is not limited to:
 - Manual breast pumps and all related equipment and supplies.
 - Hospital-grade breast pumps and all related equipment and supplies.
 - Equipment and supplies not listed in the [Covered Breastfeeding Equipment](#) section above, including but not limited to:
 - Batteries, battery-powered adaptors, and battery packs.
 - Electrical power adapters for travel.
 - Bottles which are not specific to breast pump operation. This includes the associated bottle nipples, caps, and lids.
 - Travel bags, and other similar travel or carrying accessories.
 - Breast pump cleaning supplies including soap, sprays, wipes, steam cleaning bags and other similar products.
 - Baby weight scales.
 - Garments or other products that allow hands-free pump operation.
 - Breast milk storage accessories such as ice-packs, labels, labeling lids, and other similar products. The breastmilk storage accessories exclusion does not apply to breastmilk storage bags (HCPSC code A4287).
 - Nursing bras, bra pads, breast shells, nipple shields, and other similar products.
 - Creams, ointments, and other products that relieve breastfeeding related symptoms or conditions of the breasts or nipples.

Note: Refer to the [Indications for Coverage](#) section above for covered breastfeeding equipment.

Frequently Asked Questions (FAQ)

1	Q:	If a polyp is encountered during a preventive screening colonoscopy, are future colonoscopies considered under the preventive care services benefit?
	A:	No. If a polyp is removed during a preventive screening colonoscopy, future colonoscopies would normally be considered to be diagnostic because the time intervals between future colonoscopies would be shortened.

2	Q:	Are the related therapeutic services for a preventive colonoscopy covered under the preventive care benefit?
	A:	Yes, related services integral to a colonoscopy are covered under the preventive care services benefit including: pre-operative examination, the associated facility, anesthesia, polyp removal (if necessary), pathologist and physician fees. However, the preventive benefit does not include a post-operative examination.
3	Q:	Do any preventive care services require prior-authorization?
	A:	Certain services require prior-authorization on most benefit plans. This includes, but may not be limited to: BRCA lab screening, computed tomographic colonography (virtual colonoscopy), and screening for lung cancer with low-dose computed tomography.
4	Q:	If a member in the age range of 45-75 years has a positive stool-based colorectal cancer screening test (e.g., FIT, FOBT, and fecal DNA) or direct visualization screening test (e.g., sigmoidoscopy or CT colonography), and has a follow up colonoscopy, is the colonoscopy included in the preventive care services benefit?
	A:	Yes, in this situation, the colonoscopy would be considered under the preventive care services benefit when billed in accordance with the coding in the Colorectal Cancer Screening row listed in this policy.
5	Q:	For preventive services that have a diagnosis code requirement, does the listed diagnosis code need to be the primary diagnosis on the claim?
	A:	In general, most preventive services do not require the preventive diagnosis code to be in the primary position. However, certain preventive services do require the diagnosis code to be in the primary position, which include: (1) Chemoprevention of Breast Cancer (Counseling), (2) Genetic Counseling and Evaluation for BRCA Testing, and (3) Prevention of Human Immunodeficiency Virus (HIV) Infection.
6	Q:	If a woman has an abnormal finding on a preventive screening mammography and the follow up mammogram was found to be normal, will UnitedHealthcare cover her future mammograms under the preventive care services benefit?
	A:	Yes, if the member was returned to normal mammography screening protocol, her future mammography screenings would be considered under the preventive care services benefit.
7	Q:	If a member had elevated cholesterol on a prior preventive screening, are future cholesterol tests considered under the preventive care services benefit?
	A:	Once the diagnosis has been made, further testing is considered diagnostic rather than preventive. This is true whether or not the member is receiving pharmacotherapy.
8	Q:	Are the related services for a woman's outpatient sterilization or other contraceptive procedure covered under the preventive care benefit?
	A:	Related services for a woman's outpatient sterilization or other contraceptive procedure are covered under the preventive care services benefit. This includes associated implantable devices, facility fee, anesthesia, and surgeon/physician fees. If a woman is admitted to an inpatient facility for another reason (e.g., maternity/delivery), and has a sterilization or other contraceptive procedure performed during that admission, the sterilization or other contraceptive procedure fees are covered under the preventive care services benefit. This includes associated sterilization/contraception surgical fees, device fees, anesthesia, pathology, and physician fees. However, the facility fees are not covered under the preventive care benefit since the sterilization or other contraceptive procedure is incidental to, and is not the primary reason, for the inpatient admission.
9	Q:	Are blood draws/venipunctures included in the preventive care benefit?
	A:	Yes, blood draws/venipunctures are considered under the preventive benefit if billed for a covered preventive lab service that requires a blood draw.
10	Q:	Is a newly-combined vaccine (a vaccine with several individual vaccines combined into one) covered under preventive care benefits?
	A:	A new vaccine that is pending ACIP recommendations, but is a combination of previously approved individual components, may be eligible under the preventive care benefit.
11	Q:	Are preventive care services affected by other policies?
	A:	Yes, including for example, the Reimbursement Policy titled Preventive Medicine and Screening Policy describes situations which may affect reimbursement of preventive care services.

12	Q:	Are travel vaccines covered under preventive care benefits?
	A:	Benefits for preventive care services include vaccines for routine use in children, adolescents and adults that have in effect a recommendation from ACIP with respect to the individual involved. Vaccines that are specific to travel (e.g., typhoid, yellow fever, cholera, plague, and Japanese encephalitis virus) are excluded from the preventive care services benefit.
13	Q:	Does the preventive care services benefit include prescription or over the counter (OTC) items?
	A:	Refer to the plan's pharmacy benefit plan administrator for details on prescription medications and OTCs available under the plan's preventive benefit.

Definitions

The following definitions may not apply to all plans. Refer to the member specific benefit plan document for applicable definitions.

Modifier 33: Preventive service; when the primary purpose of the service is the delivery of an evidence-based service in accordance with a US Preventive Services Task Force A or B rating in effect and other preventive services identified in preventive services mandates (legislative or regulatory), the service may be identified by adding 33 to the procedure. For separately reported services specifically identified as preventive, the modifier should not be used.

Note: UnitedHealthcare considers the procedures and diagnostic codes and Preventive Benefit Instructions listed in the table below in determining whether preventive care benefits apply. While Modifier 33 may be reported, it is not used in making preventive care benefit determinations.

Acronyms

Throughout this document the following acronyms are used:

- USPSTF: United States Preventive Services Task Force
- PPACA: Patient Protection and Affordable Care Act of 2010
- ACIP: Advisory Committee on Immunization Practices
- HRSA: Health Resources and Services Administration

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply. CPT® is a registered trademark of the American Medical Association.

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p><i>Abdominal Aortic Aneurysm Screening</i></p> <p>USPSTF Rating (Dec. 2019): B The USPSTF recommends 1-time screening for abdominal aortic aneurysm (AAA) with ultrasonography in men aged 65-75 years who have ever smoked.</p>	<p>Procedure Code(s): <i>Ultrasound Screening Study for Abdominal Aortic Aneurysm:</i> 76706</p> <p>Diagnosis Code(s): F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891</p>	<p>Age 65 through 75 (ends on 76th birthday).</p> <p>Requires at least one of the diagnosis codes listed in this row.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<i>Bacteriuria Screening</i> USPSTF Rating (Sept. 2019): A The USPSTF recommends screening for asymptomatic bacteriuria using urine culture in pregnant persons.	Procedure Code(s): 81007, 87086, 87088 Diagnosis Code(s): Pregnancy Diagnosis Codes	Requires a Pregnancy Diagnosis Code .
<i>Chlamydia Infection Screening</i> USPSTF Rating (Sept. 2021): B The USPSTF recommends screening for chlamydia in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection. Notes: <ul style="list-style-type: none"> This recommendation applies to asymptomatic, sexually active adolescents and adults, including pregnant persons. Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years. 	Procedure Code(s): <i>Chlamydia Infection Screening:</i> 86631, 86632, 87110, 87270, 87320, 87490, 87491, 87492, 87801, 87810 <i>Blood Draw:</i> 36415, 36416 Blood draw codes only apply to lab codes 86631 or 86632 Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes or <i>Screening:</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 Other: Z11.3, Z11.4, Z11.8, Z11.9, Z20.2, Z20.6, Z29.81, Z72.51, Z72.52, Z72.53	<i>Chlamydia Infection Screening:</i> Requires a Pregnancy Diagnosis Code or one of the Screening diagnosis codes listed in this row. <i>Blood Draw:</i> Required to be billed with 86631 or 86632 and <ul style="list-style-type: none"> One of the Screening diagnosis codes listed in this row or With a Pregnancy Diagnosis Code.
<i>Gonorrhea Screening</i> USPSTF Rating (Sept. 2021): B The USPSTF recommends screening for gonorrhea in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection. Notes: <ul style="list-style-type: none"> This recommendation applies to asymptomatic, sexually active adolescents and adults including pregnant persons. Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years. 	Procedure Code(s): 87590, 87591, 87592, 87801, 87850 Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes or <i>Screening:</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 Other: Z11.3, Z11.4, Z11.9, Z20.2, Z20.6, Z29.81, Z72.51, Z72.52, Z72.53	Requires either a Pregnancy Diagnosis Code or one of the Screening diagnosis codes listed in this row.

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p>Hepatitis B Virus Infection Screening</p> <p><i>Pregnant Women:</i> USPSTF Rating (July 2019): A The USPSTF recommends screening for hepatitis B virus (HBV) infection in pregnant women at their first prenatal visit.</p> <p><i>Adolescents and Adults at Increased Risk for Infection:</i> USPSTF Rating (Dec. 2020): B The USPSTF recommends screening for hepatitis B virus (HBV) infection in persons at high risk for infection.</p> <p>Bright Futures (July 2022): Bright Futures recommends screening between the ages 0-21 years (perform risk assessment for hepatitis B virus (HBV) infection).</p>	<p>Procedure Code(s): <i>Hepatitis B Virus Infection Screening:</i> 86704, 86706, 87340, 87341, 87467, G0499</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes or <i>Screening:</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 Other: Z11.3, Z11.4, Z20.2, Z20.6, Z11.59, Z29.81, Z57.8, Z72.51, Z72.52, Z72.53</p>	<p><i>Hepatitis B Virus Infection Screening:</i> Requires a Pregnancy Diagnosis Code or one of the Screening diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Requires one of the listed Hepatitis B Virus Infection Screening procedure codes listed in this row and</p> <ul style="list-style-type: none"> • A Pregnancy Diagnosis Code or • One of the Screening diagnosis codes listed in this row.
<p>Hepatitis C Virus Infection Screening</p> <p>USPSTF Rating (March 2020): B The USPSTF recommends screening for hepatitis C virus infection in adults aged 18-79 years.</p> <p>Bright Futures (March 2021) Bright Futures recommends screening all individuals ages 18 to 79 years at least once for hepatitis C virus infection (HCV).</p>	<p>Procedure Code(s): <i>Hepatitis C Virus Infection Screening:</i> 86803, 86804, G0472</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p><i>Hepatitis C Virus Infection Screening:</i> Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p><i>Blood Draw:</i> Requires one of the Hepatitis C Virus Infection Screening procedure codes listed in this row</p>
<p>HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults</p> <p>USPSTF Rating (June 2019): A The USPSTF recommends that clinicians screen for HIV infection in:</p> <ul style="list-style-type: none"> • Adolescents and adults aged 15-65 years. Younger 	<p>Procedure Code(s): <i>HIV (Human Immunodeficiency Virus) Screening:</i> 86689, 86701, 86702, 86703, 87389, 87390, 87391, 87534, 87535, 87536, 87537, 87538, 87539, 87806, G0432, G0433, G0435, G0475, S3645</p> <p><i>Blood Draw:</i> 36415, 36416</p>	<p>No age limits.</p> <p><i>HIV – Human Immunodeficiency Virus – Screening:</i> Requires a Pregnancy Diagnosis Code or one of the Screening diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Requires both of the following:</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p>adolescents and older adults who are at increased risk of infection should also be screened.</p> <ul style="list-style-type: none"> All pregnant persons, including those who present in labor or at delivery whose HIV status is unknown. <p>Note: Bright Futures recommends HIV screening lab work be conducted at least once between ages 15-21 years. Also recommended anytime between ages 11-14 years, when a risk assessment is positive. And after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently if at high risk.</p>	<p>Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes or <i>Screening:</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129, Other: Z11.3, Z11.4, Z11.59, Z11.9, Z20.2, Z20.6, Z22.6, Z22.8, Z22.9, Z29.81, Z72.51, Z72.52, Z72.53 Also see Expanded Women's Preventive Health section.</p>	<ul style="list-style-type: none"> One of the listed HIV Screening procedure codes listed in this row and One of the Screening diagnosis codes listed in this row or a Pregnancy Diagnosis Code.
<p><i>RH Incompatibility Screening</i></p> <p>USPSTF Rating (Feb. 2004): A Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.</p> <p>USPSTF Rating (Feb. 2004): B Repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.</p>	<p>Procedure Code(s): <i>RH Incompatibility Screening:</i> 86850, 86901</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Pregnancy Diagnosis Codes</p>	<p><i>RH Incompatibility Screening:</i> Requires a Pregnancy Diagnosis Code.</p> <p><i>Blood Draw:</i> Required to be billed with 86850 or 86901 and with a Pregnancy Diagnosis Code.</p>
<p><i>Syphilis Screening</i></p> <p><i>Non-Pregnant Adolescents and Adults at Increased Risk:</i></p> <p>USPSTF Rating (Sept. 2022): A The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection (asymptomatic, nonpregnant adolescents and adults who are at increased risk for syphilis infection).</p> <p><i>Pregnant Women:</i></p> <p>USPSTF Rating (Sept. 2018): A</p>	<p>Procedure Code(s): <i>Syphilis Screening:</i> 0064U, 0065U, 0210U, 86592, 86593, 86780</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): <i>Pregnancy:</i> Pregnancy Diagnosis Codes or <i>Screening:</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129</p>	<p><i>Syphilis Screening:</i> Requires a Pregnancy Diagnosis Code or one of the Screening diagnosis code listed in this row.</p> <p><i>Blood Draw:</i> Requires both of the following:</p> <ul style="list-style-type: none"> One of the listed Syphilis Screening procedure codes listed in this row and One of the Screening diagnosis codes listed in this row or a Pregnancy Diagnosis Code.

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
The USPSTF recommends early screening for syphilis infection in all pregnant women. Note: Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years.	Other: Z11.2, Z11.3, Z11.4, Z11.9, Z20.2, Z20.6, Z29.81, Z72.51, Z72.52, Z72.53	
Genetic Counseling and Evaluation for BRCA Testing; and BRCA Lab Screening USPSTF Rating (Aug. 2019): B The USPSTF recommends that primary care clinicians assess women with a personal or family history of breast, ovarian, tubal, or peritoneal cancer or who have an ancestry associated with breast cancer susceptibility 1 and 2 (BRCA1/2) gene mutations with an appropriate brief familial risk assessment tool. Women with a positive result on the risk assessment tool should receive genetic counseling and, if indicated after counseling, genetic testing. Refer to the Medical Policy titled Genetic Testing for Hereditary Cancer .	Genetic Counseling and Evaluation Procedure Code(s): <i>Medical Genetics and Genetic Counseling Services:</i> 96041, S0265 <i>Evaluation and Management (Office Visits):</i> 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, 99385, 99386, 99387, 99395, 99396, 99397, G0463 Diagnosis Code(s): Z15.01, Z15.02, Z80.3, Z80.41, Z85.3, Z85.43	Genetic Counseling and Evaluation *Medical Necessity plans require genetic counseling before BRCA Lab Screening. Requires one of the Genetic Counseling and Evaluation diagnosis codes listed in this row in the primary position.
Refer to the Medical Policy titled Genetic Testing for Hereditary Cancer .	BRCA Lab Screening Procedure Code(s): 81162, 81163, 81164, 81165, 81166, 81167, 81212, 81215, 81216, 81217 <i>Blood Draw:</i> 36415, 36416 Diagnosis Code(s): <i>Family History or Personal History of breast cancer and/or ovarian cancer:</i> Z15.01, Z15.02, Z80.3, Z80.41, Z85.3, Z85.43	BRCA Lab Screening *Prior authorization requirements apply to BRCA lab screening. Applies to age 18+ when billed with one of the BRCA Lab Screening diagnosis codes listed in this row. <i>Blood Draw:</i> Requires one of the BRCA Lab Screening procedure codes listed in this row and one of the BRCA Lab Screening diagnosis codes listed in this row.
Screening for Pre-Diabetes and Type 2 Diabetes USPSTF Rating (Aug. 2021): B The USPSTF recommends screening for prediabetes and type 2 diabetes in adults aged 35 to 70 years who have overweight or obesity. Clinicians should offer or	Pre-Diabetes Preventive Interventions Procedure Code(s): <i>Medical Nutrition Therapy or Counseling:</i> 97802, 97803, 97804, G0270, G0271, S9470 <i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404	Pre-Diabetes Preventive Interventions Limited to age 35-70 years (ends on 71 st birthday). Requires diagnosis code R73.03.

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
refer patients with prediabetes to effective preventive interventions. Refer to Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors: Behavioral Counseling Interventions for intensive behavioral counseling interventions.	<i>Behavioral Counseling or Therapy:</i> 0403T, G0447, G0473, G9886 Diagnosis Code(s): R73.03 (prediabetes)	
For additional diabetes screening benefits, also see the <i>Expanded Women's Preventive Health</i> section for Screening for Diabetes in Pregnancy and Screening for Diabetes After Pregnancy .	Diabetes Screening Procedure Code(s): <i>Diabetes Screening:</i> 82947, 82948, 82950, 82951, 82952, 83036 <i>Blood Draw:</i> 36415, 36416 Diagnosis Code(s): <i>Required Diagnosis Codes (requires at least one):</i> Z00.00, Z00.01, Z13.1 And one of the following additional diagnosis codes as follows: <i>Additional Diagnosis Codes (requires at least one):</i> <i>Overweight:</i> E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29 <i>Obesity:</i> E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 <i>Body Mass Index 30.0 – 39.9:</i> Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39 <i>Body Mass Index 40.0 and Over:</i> Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 See the <i>Expanded Women's Preventive Health</i> section for Screening for Diabetes in Pregnancy and Screening for Diabetes After Pregnancy .	Diabetes Screening Limited to age 35-70 years (ends on 71 st birthday). <i>Diabetes Screening:</i> Requires one of the Required Diagnosis Codes listed in this row and one of the listed Additional Diagnosis Codes in this row. <i>Blood Draw:</i> Requires all of the following: <ul style="list-style-type: none"> ● One of the listed Diabetes Screening procedure codes listed in this row and ● One of the listed Required Diagnosis Codes and ● One of the listed Additional Diagnosis Codes. Preventive Benefit Does Not Apply: If a Diabetes Diagnosis Code is present in any position, the preventive benefit does not apply; see the Diabetes Diagnosis Code List .

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p><i>Gestational Diabetes Screening</i></p> <p>USPSTF Rating (Aug. 2021): B The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant persons at 24 weeks of gestation or after.</p> <p>For additional diabetes screening benefits, also see the Screening for Pre-Diabetes and Type 2 Diabetes row. Also see the <i>Expanded Women's Preventive Health</i> section for Screening for Diabetes in Pregnancy and Screening for Diabetes After Pregnancy.</p>	<p>See the <i>Expanded Women's Preventive Health</i> section for Screening for Diabetes in Pregnancy codes.</p>	<p>See the <i>Expanded Women's Preventive Health</i> section for Screening for Diabetes in Pregnancy preventive benefit instructions.</p> <p>Note: This benefit applies regardless of the gestational week.</p>
<p><i>Screening Mammography</i></p> <p>USPSTF Rating (2002): B The USPSTF recommends screening mammography, with or without clinical breast examination (CBE), every 1-2 years for women aged 40 and older.</p> <p>Also refer to the Medical Policy titled Breast Imaging for Screening and Diagnosing Cancer.</p> <p>Also see the Breast Cancer Screening for Average-Risk Women recommendation in the <i>Expanded Women's Preventive Health</i> section.</p>	<p>Procedure Code(s): 77063, 77067</p> <p>Revenue Code: 0403</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>No age limits.</p> <p>Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>Note: This benefit only applies to screening mammography.</p>
<p><i>Cervical Cancer Screening</i></p> <p>USPSTF Rating (Aug. 2018): A <i>Age 21-29 years:</i> The USPSTF recommends screening for cervical cancer every</p>	<p>Human Papillomavirus DNA Testing (HPV)</p> <p>Procedure Code(s): 0502U, 87624, 87625, 87626, G0476</p> <p>Diagnosis Code(s): Z00.00, Z00.01, Z01.411, Z01.419, Z11.51, Z12.4</p>	<p>Human Papillomavirus DNA Testing (HPV)</p> <p>Age 30 years and up. Requires one of the diagnosis codes listed in this row.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p>3 years with cervical cytology alone in women aged 21-29 years.</p> <p><u>Age 30-65 years:</u> For women aged 30 to 65 years, the USPSTF recommends:</p> <ul style="list-style-type: none"> • Screening every 3 years with cervical cytology alone, • Every 5 years with high-risk human papillomavirus (hrHPV) testing alone, or • Every 5 years with hrHPV testing in combination with cytology (co-testing). <p>Bright Futures, March 2014: Adolescents should no longer be routinely screened for cervical dysplasia until age 21.</p> <p>Also see Screening for Cervical Cancer in the <i>Expanded Women's Preventive Health</i> section.</p>	<p>Cervical Cytology (Pap Test) Code Group 1 Procedure Code(s): G0101, G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, Q0091, P3000, P3001</p> <p>Code Group 1 Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.</p> <p>Code Group 2 Procedure Code(s): 88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88155, 88164, 88165, 88166, 88167, 88174, 88175</p> <p>Code Group 2 Diagnosis Code(s): Z00.00, Z00.01, Z01.411, Z01.419, Z12.4</p>	<p>Cervical Cytology (Pap Test) Code Group 1: Limited to age 21-65 years (ends on 66th birthday). Does not have diagnosis code requirements for preventive benefits to apply.</p> <p>Code Group 2: Limited to age 21-65 years (ends on 66th birthday). Requires one of the Code Group 2 diagnosis codes listed in this row.</p>
<p><i>Statin Use for the Primary Prevention of Cardiovascular Disease in Adults - Cholesterol Screening (Lipid Disorders Screening)</i></p> <p>USPSTF Rating (August 2022): B The USPSTF recommends that clinicians prescribe a statin for the primary prevention of CVD for adults aged 40 to 75 years who have 1 or more CVD risk factors (i.e. dyslipidemia, diabetes, hypertension, or smoking) and an estimated 10-year risk of a cardiovascular event of 10% or greater.</p> <p>Notes:</p> <ul style="list-style-type: none"> • For statin medications benefits, refer to the pharmacy plan administrator. • See Dyslipidemia Screening (Bright Futures) for recommendations for children. 	<p>Procedure Code(s): <i>Cholesterol Screening:</i> 80061, 82465, 83718, 83719, 83721, 83722, 84478</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p><i>ASCVD Risk Assessment and Risk Management Services:</i> G0537, G0538</p> <p>Diagnosis Code(s): Z00.00, Z00.01, Z13.220</p>	<p><i>Cholesterol Screening:</i> Ages 40-75 years (ends on 76th birthday). Requires one of the diagnosis codes listed in this row for CPT codes 80061, 82465, 83718, 83719, 83721, 83722, and 84478.</p> <p><i>Blood Draw:</i> Ages 40-75 years (ends on 76th birthday): Requires one of the listed Cholesterol Screening procedure codes and one of the Diagnosis Codes listed in this row.</p> <p><i>ASCVD Risk Assessment and Management Services:</i> Ages 40-75 years (ends on 76th birthday). The diagnosis codes listed in this row are not required for HCPCS codes G0537 and G0538.</p> <p>Preventive Benefit Does Not Apply: For all ages above, if any of the following lipid disorders diagnosis codes are present in any position, the preventive benefit does not apply: E71.30, E75.5, E78.00, E78.01, E78.2,</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p>Colorectal Cancer Screening</p> <p>USPSTF Rating (May 2021): B The USPSTF recommends screening for colorectal cancer in adults aged 45 to 49 years.</p> <p>USPSTF Rating (May 2021): A The USPSTF recommends screening for colorectal cancer in all adults aged 50 to 75 years.</p> <p>Also refer to the Medical Policies titled Outpatient Surgical Procedures – Site of Service; Screening Colonoscopy Procedures – Site of Service; and Magnetic Resonance Imaging (MRI) and Computed Tomography (CT) Scan – Site of Service.</p> <p>Also see the Frequently Asked Questions section.</p>	<p>Colonoscopy Procedure Code(s): <i>Preventive Colonoscopy:</i> G0105, G0121</p> <p><i>Preventive Colonoscopy When Billed with Certain Codes (see Preventive Benefit Instructions to the right):</i> 44388*, 44389*, 44392*, 44394*, 45378*, 45380*, 45381*, 45384*, 45385*, 45388*</p> <p>Diagnosis Code(s): <i>Applies to Procedure Codes with asterisk(*) above:</i> Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79</p> <p>Note: Also see the Colonoscopy Pre-Op Consultation row below.</p>	<p>Colonoscopy Age Limits: 45-75 years (ends on 76th birthday).</p> <p>HCPCS codes G0105 and G0121 do not have diagnosis code requirements for preventive benefits to apply. Codes with an asterisk(*) are preventive when:</p> <ul style="list-style-type: none"> Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or Billed in addition to HCPCS codes G0104, G0105, G0106, G0120, G0121, G0122, G0328, or S0285.
	<p>Sigmoidoscopy Procedure Code(s): <i>Preventive Sigmoidoscopy:</i> G0104</p> <p><i>Preventive Sigmoidoscopy When Billed with Certain Codes (see Preventive Benefit Instructions to the right):</i> 45330*, 45331*, 45333*, 45338*, 45346*</p> <p>Diagnosis Code(s): <i>Applies to Procedure Codes with asterisk(*) above:</i> Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79,</p>	<p>Sigmoidoscopy Age Limits: 45-75 years (ends on 76th birthday).</p> <p>HCPCS code G0104 does not have diagnosis code requirements for preventive benefits to apply.</p> <p>CPT codes with an asterisk(*) are preventive when:</p> <ul style="list-style-type: none"> Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or Billed in addition to HCPCS codes G0104, G0105, G0106, G0120, G0121, G0122, G0328, or S0285.
	<p>Barium Enema Procedure Code(s): G0106, G0120, G0122</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.</p>	<p>Barium Enema Age Limits: 45-75 years (ends on 76th birthday).</p> <p>Codes G0106, G0120, and G0122 do not have diagnosis code requirements for preventive benefits to apply.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
	<p>Pathology and Anesthesia (for Colonoscopy or Sigmoidoscopy)</p> <p>Procedure Code(s): <i>Pathology:</i> 88304, 88305</p> <p><i>Anesthesia:</i> 00812, 99152, 99153, 99156, 99157, G0500</p> <p>Diagnosis Code(s): <i>Applies to the Pathology and Anesthesia codes listed above:</i> Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79</p>	<p>Pathology and Anesthesia (for Colonoscopy or Sigmoidoscopy)</p> <p>Age Limits: 45-75 years (ends on 76th birthday).</p> <p>Requires both of the following:</p> <ul style="list-style-type: none"> One of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); and One of the procedure codes listed in the Colonoscopy row, or the Sigmoidoscopy row. <p>Note: Preventive benefits apply when the surgeon's claim is preventive.</p>
	<p>Fecal Occult Blood Testing (FOBT) and Fecal Immunochemical Test (FIT)</p> <p>Procedure Code(s): <i>Preventive:</i> G0328</p> <p><i>Preventive When Billed with Certain Codes (see Preventive Benefit Instructions to the right):</i> 82270*, 82274*</p> <p>Diagnosis Code(s): <i>Applies to Procedure Codes with asterisk(*) above:</i> Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79</p>	<p>Fecal Occult Blood Testing (FOBT) and Fecal Immunochemical Test (FIT)</p> <p>Age Limits: 45-75 years (ends on 76th birthday).</p> <p>HCPCS code G0328 does not have diagnosis code requirements for preventive benefits to apply.</p> <p>CPT codes with an asterisk(*) are preventive when:</p> <ul style="list-style-type: none"> Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or Billed in addition to HCPCS codes G0104, G0105, G0106, G0120, G0121, G0122, G0328, or S0285.
	<p>Fecal DNA</p> <p>Procedure Code(s): 0464U, 81528</p> <p>Diagnosis Code(s): CPT code 81528 does not have diagnosis code requirements for preventive benefits to apply.</p>	<p>Fecal DNA</p> <p>Age Limits: 45-75 years (ends on 76th birthday).</p> <p>Benefit is limited to once every 3 years.</p> <p>CPT codes 0464U and 81528 do not have diagnosis code requirements for preventive benefits to apply.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
	<p>Pre-Op Consultation</p> <p>Procedure Code(s): <i>Preventive:</i> S0285</p> <p><i>Preventive when billed with one of the diagnosis codes listed in this row:</i> 99202*, 99203*, 99204*, 99205*, 99211*, 99212*, 99213*, 99214*, 99215*, 99242*, 99243*, 99244*, 99245*, 99417*</p> <p>Diagnosis Code(s): <i>Applies to Procedure Codes with asterisk(*) above:</i> Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79</p> <p>Note: For additional information on the reimbursement of consultation codes 99242-99245, refer to the Reimbursement Policy titled Consultation Services.</p>	<p>Pre-Op Consultation</p> <p>Age Limits: 45-75 years (ends on 76th birthday).</p> <p>HCPCS code S0285 does not have diagnosis code requirements for preventive benefits to apply.</p> <p>Codes with an asterisk(*) are preventive when billed with one of the diagnosis codes listed in this row (Z12.10, Z12.11, Z12.12, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79).</p>
	<p>Computed Tomographic Colonography (Virtual Colonoscopy)</p> <p>Procedure Code(s): 74263</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.</p>	<p>Computed Tomographic Colonography (Virtual Colonoscopy)</p> <p>Age Limits: 45-75 years (ends on 76th birthday).</p> <p>Does not have diagnosis code requirements for preventive benefit to apply.</p> <p>Prior authorization requirements may apply, depending on plan.</p>
<p>Wellness Examinations (well-baby, well-child, well-adult)</p> <p>USPSTF Rating: None UnitedHealthcare supports AAP and AAFP age and frequency guidelines.</p> <p>HRSA Requirements: The Wellness Examinations codes in this row include the following HRSA requirements for women, where applicable:</p>	<p>Procedure Code(s): <i>Medicare Wellness Exams:</i> G0402, G0438, G0439</p> <p><i>STIs Behavioral Counseling:</i> G0445</p> <p><i>Annual Gynecological Exams:</i> S0610, S0612, S0613</p> <p><i>Pelvic Examination (add-on code):</i> 99459</p>	<p>Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>HCPCS code G0445 is limited to twice per year.</p> <p>HCPCS code G0296 is limited to age 50 to 80 years (ends on 81st birthday).</p> <p><i>Pelvic Examination add-on code 99459:</i> Preventive care services benefits may apply to CPT code 99459 when the related evaluation and management (office visit) code is applied to the</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<ul style="list-style-type: none"> • Breastfeeding support, counseling, and education • Contraceptive methods and sterilizations (counseling and follow-up care) • Screening and counseling for interpersonal domestic violence • Screening for human immunodeficiency virus infection (HIV); education and risk assessment • Counseling for sexually transmitted infections (STIs) • Well-woman preventive visits • Screening for urinary incontinence • Obesity prevention in midlife women (counseling) 	<p><i>Preventive Medicine Services (Evaluation and Management):</i> 99381, 99382, 99383, 99384, 99385, 99386, 99387 99391, 99392, 99393, 99394, 99395, 99396, 99397</p> <p><i>Preventive Medicine, Individual Counseling:</i> 99401, 99402, 99403, 99404</p> <p><i>Preventive Medicine, Group Counseling:</i> 99411, 99412</p> <p><i>Newborn Care (evaluation and management):</i> 99461</p> <p><i>Counseling Visit (to discuss the need for Lung Cancer Screening (LDCT) using Low Dose CT Scan):</i> G0296</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>Also see the Expanded Women's Preventive Health section.</p>	<p>preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.</p>
<p><i>Newborn Screenings</i> All newborns</p> <p>USPSTF Rating (March 2008): A Hypothyroidism Screening: Screening for congenital hypothyroidism in newborns.</p> <p>USPSTF Rating (March 2008): A Phenylketonuria Screening: Screening for phenylketonuria (PKU) in newborns.</p> <p>USPSTF Rating (Sept. 2007): A Sickle Cell Screening: Screening for sickle cell disease in newborns.</p> <p>Note: For Bright Futures hearing screening, see Hearing Tests (Bright Futures).</p>	<p>Procedure Code(s):</p> <p><i>Hypothyroidism Screening:</i> 84437, 84443</p> <p><i>Phenylketonuria Screening:</i> 84030, S3620</p> <p><i>Sickle Cell Screening:</i> 83020, 83021, 83030, 83033, 83051, S3850</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p><i>Newborn Screenings:</i> Age 0-90 days. Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p><i>Blood Draw:</i> Age 0-90 days, requires one of the listed Hypothyroidism Screening, Phenylketonuria Screening, or Sickle Cell Screening procedure codes.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
Metabolic Screening Panel (Newborns)	Procedure Code(s): <i>Metabolic Screening Panel:</i> 82017, 82136, 82261, 82775, 83020, 83498, 83516, 84030, 84437, 84443, S3620 <i>Blood Draw:</i> 36415, 36416 Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.	<i>Metabolic Screening Panel:</i> Age 0-90 days. Does not have diagnosis code requirements for the preventive benefit to apply. <i>Blood Draw:</i> Age 0-90 days. Requires one of the listed Metabolic Screening Panel procedure codes listed in this row.
Osteoporosis Screening USPSTF Rating (June 2018): B Women 65 and older: The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in women 65 years and older. USPSTF Rating (June 2018): B Postmenopausal women younger than 65 years at increased risk of osteoporosis: The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in postmenopausal women younger than 65 years who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool.	Procedure Code(s): 76977, 77080, 77081, G0130 Diagnosis Code(s): Z00.00, Z00.01, Z13.820, Z82.62	Requires one of the diagnosis codes listed in this row.
Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults USPSTF Rating (Nov. 2018): B The USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant women, and providing persons	Procedure Code(s): <i>Alcohol or Drug Use Screening:</i> 99408, 99409 <i>Annual Alcohol Screening:</i> G0442 <i>Brief Counseling for Alcohol:</i> G0443 Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.	Does not have diagnosis code requirements for preventive benefits to apply.

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p>engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy alcohol use.</p> <p>Bright Futures (April 2017): Bright Futures recommends alcohol or drug use assessments from age 11-21 years.</p> <p>Also see rows: Unhealthy Drug Use Screening (Adults); and Tobacco, Alcohol, or Drug Use Assessment (Bright Futures).</p>		
<p><i>Unhealthy Drug Use Screening (Adults)</i></p> <p>USPSTF Rating (June 2020): B The USPSTF recommends screening by asking questions about unhealthy drug use in adults age 18 years or older. Screening should be implemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers to asking questions about unhealthy drug use, not testing biological specimens.)</p> <p>Bright Futures (April 2017): Bright Futures recommends alcohol or drug use assessments from age 11-21 years.</p> <p>Also see rows: Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults; and Tobacco, Alcohol, or Drug Use Assessment (Bright Futures).</p>	<p>Procedure Code(s): <i>Alcohol or Drug Use Screening:</i> 99408, 99409</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for preventive benefit to apply.</p>	<p>Does not have diagnosis code requirements for preventive benefits to apply.</p>
<p><i>High Blood Pressure in Adults – Screening</i></p> <p>USPSTF Rating (April 2021): A The USPSTF recommends screening for hypertension in adults 18 years or older with office blood pressure measurement.</p>	<p>Blood Pressure Measurement in a Clinical Setting N/A</p>	<p>Blood Pressure Measurement in a Clinical Setting This service is included in a preventive care wellness examination.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
The USPSTF recommends obtaining blood pressure measurements outside of the clinical setting for diagnostic confirmation before starting treatment.	Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting) Procedure Code(s): <i>Ambulatory Blood Pressure Measurement:</i> 93784, 93786, 93788 or 93790 Diagnosis Code(s): <i>Abnormal Blood-Pressure Reading Without Diagnosis of Hypertension:</i> R03.0	Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting) Age 18 years and older. Requires the diagnosis code listed in this row.
Breast Cancer: Medication Use to Reduce Risk USPSTF Rating (Sept. 2019): B The USPSTF recommends that clinicians offer to prescribe risk-reducing medications, such as tamoxifen, raloxifene, or aromatase inhibitors, to women who are at increased risk for breast cancer and at low risk for adverse medication effects.	Procedure Code(s): <i>Evaluation and Management (Office Visits):</i> 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99385, 99386, 99387, 99395, 99396, 99397, 99417, G0463 Diagnosis Code(s): Z80.3, Z80.41, Z15.01, Z15.02	Requires one of the diagnosis codes listed in this row in the primary position.
Primary Care Interventions to Promote Breastfeeding USPSTF Rating (Oct. 2016): B The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.	N/A Also see the Expanded Women's Preventive Health section	Included in primary care or OB/GYN office visits.
Depression in Adults (Screening) USPSTF Rating (June 2023): B The USPSTF recommends screening for depression in the adult population, including pregnant and postpartum persons, as well as older adults. Bright Futures (February 2017): Maternal Depression Screening: Routine screening for postpartum depression should be integrated	Procedure Code(s): 96127, 96161, G0136, G0444 Diagnosis Code(s): Required for 96127 Only: <i>Encounter for Screening for Depression:</i> Z13.31, Z13.32	Requires one of the diagnosis codes listed in this row, for 96127. The diagnosis codes listed in this row are not required, for G0136, G0444, and 96161.

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
into well-child visits at 1, 2, 4, and 6 months of age. Also see the rows for Screening for Anxiety (HRSA); Anxiety Disorders in Adults Screening (USPSTF); Depression in Children and Adolescents (Screening) (USPSTF); Perinatal Depression – Preventive Interventions (Counseling) (USPSTF); and Depression and Suicide Risk Screening (Bright Futures).		
<p><i>Depression in Children and Adolescents (Screening)</i></p> <p>USPSTF Rating (October 2022): B The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12-18 years.</p> <p>Bright Futures (February 2017): Maternal Depression Screening: Routine screening for postpartum depression should be integrated into well-child visits at 1, 2, 4, and 6 months of age.</p> <p>Note: The Bright Futures Periodicity Schedule recommends depression screening begin at age 12-21 years.</p> <p>Also see the rows for Anxiety Disorders in Adults Screening (USPSTF); Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Perinatal Depression – Preventive Interventions (Counseling) (USPSTF); and Depression and Suicide Risk Screening (Bright Futures).</p>	<p>Procedure Code(s): 96127, 96161, G0136, G0444</p> <p>Diagnosis Code(s): Required for 96127 Only: <i>Encounter for Screening for Depression:</i> Z13.31, Z13.32</p>	<p>Requires one of the diagnosis codes listed in this row for 96127.</p> <p>The diagnosis codes listed in this row are not required for G0136, G0444, and 96161.</p>
<p><i>Anxiety Disorders in Adults (Screening)</i></p> <p>USPSTF Rating (June 2023): B The USPSTF recommends screening for anxiety in adults, including pregnant and postpartum</p>	<p>Procedure Code(s): 96127</p>	<p>Requires the diagnosis code listed in this row.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
persons. This applies to adults age 64 or younger. Also see the rows for Screening for Anxiety (HRSA) ; and Screening for Anxiety in Children and Adolescents (USPSTF) .	Diagnosis Code(s): <i>Encounter for Screening Examination for Other Mental Health and Behavioral Disorders:</i> Z13.39	
Screening for Anxiety in Children and Adolescents USPSTF Rating (October 2022): B The USPSTF recommends screening for anxiety in children and adolescents aged 8 to 18 years. Also see the rows for Anxiety Disorders in Adults Screening (USPSTF) ; Screening for Anxiety (HRSA) ; Screening for Depression in Adults (USPSTF) ; Perinatal Depression – Preventive Interventions (Counseling) (USPSTF) ; and Depression and Suicide Risk Screening (Bright Futures) .	Procedure Code(s): 96127 Diagnosis Code(s): <i>Encounter for Screening Examination for Other Mental Health and Behavioral Disorders:</i> Z13.39	Requires the diagnosis code listed in this row.
Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors: Behavioral Counseling Interventions USPSTF Rating (Nov. 2020): B The USPSTF recommends offering or referring adults with cardiovascular disease risk factors to behavioral counseling interventions to promote a healthy diet and physical activity.	Procedure Code(s): <i>Medical Nutrition Therapy or Counseling:</i> 97802, 97803, 97804, G0270, G0271, S9470 <i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404 <i>Behavioral Counseling or Therapy:</i> 0403T, G0446, G0447, G0473, G9886 <i>ASCVD Risk Assessment and Risk Management Services:</i> G0537, G0538 Diagnosis Code(s): <i>Screening:</i> Z13.220 <i>Nicotine Dependence, Tobacco Use, or Family History of IHD:</i> F17.210, F17.211, F17.213, F17.218, F17.219, Z72.0, Z87.891, Z82.49	Requires one of the diagnosis codes listed in this row for CPT/HCPCS codes 0403T, 97802-97804, 99401-99404, G0270, G0271, G9886, and S9470. The diagnosis codes listed in this row are not required for HCPCS codes G0446, G0447, G0473, G0537, and G0538. HCPCS code G0446 is limited to once per year.

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
	<p><i>Overweight:</i> E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29</p> <p><i>Body Mass Index 30.0 – 39.9:</i> Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39</p> <p><i>Body Mass Index 40.0 and Over:</i> Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</p> <p><i>Impaired Fasting Glucose:</i> R73.01</p> <p><i>Metabolic Syndrome; Insulin Resistance Syndrome Type A; Other Insulin Resistance:</i> E88.810, E88.811, E88.818, E88.819</p> <p><i>Hyperlipidemia / Dyslipidemia:</i> E78.00, E78.01, E78.1, E78.2, E78.3, E78.41, E78.49, E78.5</p> <p><i>Obesity:</i> E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</p> <p><i>Essential Hypertension:</i> I10</p> <p><i>Resistant Hypertension:</i> I1A.0</p> <p><i>Secondary Hypertension:</i> I15.0, I15.1, I15.2, I15.8, I15.9, N26.2</p> <p><i>Hypertension Complicating Pregnancy, Childbirth, and the Puerperium:</i> O10.011, O10.012, O10.013, O10.019, O10.02, O10.03, O10.111, O10.112, O10.113, O10.119, O10.12, O10.13, O10.211, O10.212, O10.213, O10.219, O10.22, O10.23, O10.311, O10.312, O10.313, O10.319, O10.32, O10.33, O10.411, O10.412, O10.413, O10.419, O10.42, O10.43, O10.911, O10.912, O10.913, O10.919, O10.92,</p>	

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
	O10.93, O11.1, O11.2, O11.3, O11.4, O11.5, O11.9, O13.1, O13.2, O13.3, O13.4, O13.5, O13.9, O16.1, O16.2, O16.3, O16.4, O16.5, O16.9 <i>Urgent/Emergency/Crisis Hypertension:</i> I16.0, I16.1, I16.9 <i>Diabetes:</i> Diabetes Diagnosis Code List <i>Atherosclerosis:</i> Atherosclerosis Diagnosis Code List <i>Coronary Atherosclerosis:</i> I25.10, I25.110, I25.111, I25.112, I25.118, I25.119, I25.700, I25.701, I25.702, I25.708, I25.709, I25.710, I25.711, I25.712, I25.718, I25.719, I25.720, I25.721, I25.722, I25.728, I25.729, I25.730, I25.731, I25.732, I25.738, I25.739, I25.750, I25.751, I25.752, I25.758, I25.759, I25.760, I25.761, I25.762, I25.768, I25.769, I25.790, I25.791, I25.792, I25.798, I25.799, I25.810, I25.811, I25.812	
<p><i>Weight Loss to Prevent Obesity-Related Morbidity and Mortality in Adults: Behavioral Interventions</i></p> <p>USPSTF Rating (Sept. 2018): B The USPSTF recommends that clinicians offer or refer adults with a body mass index (BMI) of 30 or higher (calculated as weight in kilograms divided by height in meters squared) to intensive multicomponent behavioral interventions.</p>	<p>Procedure Code(s): <i>Medical Nutrition Therapy:</i> 97802, 97803, 97804, G0270, G0271, S9470 <i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404 <i>Behavioral Counseling or Therapy:</i> 0403T, G0446, G0447, G0473, G9886 Diagnosis Code(s): <i>Body Mass Index 30.0-39.9:</i> Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39 <i>Body Mass Index 40.0 and over:</i> Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</p>	<p>Requires one of the diagnosis codes listed in this row for CPT/HCPCS codes 0403T, 97802-97804, 99401-99404, G0270, G0271, G9886, and S9470.</p> <p>HCPCS code G0446 is limited to once per year.</p> <p>The diagnosis codes listed in this row are not required for HCPCS codes G0446, G0447, and G0473.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
	Obesity: E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82	
<i>High Body Mass Index in Children and Adolescents</i> USPSTF Rating (June 2024): B The USPSTF recommends that clinicians provide or refer children and adolescents 6 years or older with a high body mass index (BMI) (\geq 95 th percentile for age and sex) to comprehensive, intensive behavioral interventions. See the <i>Practice Considerations</i> section of the published USPSTF recommendation or more information about behavioral interventions.	Procedure Code(s): <i>Medical Nutrition Therapy:</i> 97802, 97803, 97804, G0270, G0271, S9470 <i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404 <i>Behavioral Counseling or Therapy:</i> 0403T, G0446, G0447, G0473, G9886 Also see the codes in the Wellness Examinations row above. Diagnosis Code(s): <i>Obesity:</i> E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82 <i>Pediatric BMI of 120% or more of the 95th percentile for age:</i> Z68.54, Z68.55, Z68.56	Requires one of the diagnosis codes listed in this row for CPT/ HCPCS codes 0403T, 97802-97804, 99401-99404, G0270, G0271, G9886, and S9470. HCPCS code G0446 is limited to once per year. The diagnosis codes listed in this row are not required for HCPCS codes G0446, G0447, and G0473.
<i>Healthy Weight and Weight Gain During Pregnancy: Behavioral Counseling Interventions</i> USPSTF Rating (May 2021): B The USPSTF recommends that clinicians offer pregnant persons effective behavioral counseling interventions aimed at promoting healthy weight gain and preventing excess gestational weight gain in pregnancy.	Procedure Code(s): <i>Medical Nutrition Therapy:</i> 97802, 97803, 97804, G0270, G0271, S9470 <i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404 <i>Behavioral Counseling or Therapy:</i> G0447, G0473 Diagnosis Code(s): Pregnancy Diagnosis Codes	Requires one of the diagnosis codes listed in this row.

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p><i>Behavioral Counseling to Prevent Sexually Transmitted Infections</i></p> <p>USPSTF Rating (Aug. 2020): B The USPSTF recommends behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections (STIs).</p>	<p>Procedure Code(s): <i>STIs Behavioral Counseling:</i> G0445</p> <p><i>Preventive Medicine Individual Counseling</i> 99401, 99402, 99403, 99404</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>HCPCS code G0445 is limited to twice per year.</p>
<p><i>Interventions for Tobacco Smoking Cessation in Adults, including Pregnant Persons</i></p> <p>USPSTF Rating (Jan. 2021): A Pregnant Persons (A): The USPSTF recommends that clinicians ask all pregnant persons about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant persons who use tobacco.</p> <p>Nonpregnant Adults (A): The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and US Food and Drug Administration (FDA)–approved pharmacotherapy for cessation to nonpregnant adults who use tobacco.</p> <p>Note: Refer to the plan's pharmacy benefit plan administrator for details on prescription medications available under the plan's preventive benefit.</p> <p>Also see rows: Unhealthy Drug Use Screening (Adults); and Tobacco, Alcohol, or Drug Use Assessment (Bright Futures).</p>	<p>Procedure Code(s): <i>Behavioral Interventions:</i> 99406, 99407</p> <p><i>Preventive Medicine, Individual Counseling:</i> 99401, 99402, 99403, 99404</p> <p>Also see the codes in the Wellness Examinations row above.</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>Does not have diagnosis code requirements for the preventive benefit to apply.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p>Primary Care Interventions To Prevent Tobacco Use In Children And Adolescents</p> <p>USPSTF Rating (April 2013): B The USPSTF recommends that primary care clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use among school-aged children and adolescents.</p> <p>Bright Futures (April 2017): Bright Futures recommends tobacco use assessments from age 11-21 years.</p> <p>Also see rows: Unhealthy Drug Use Screening (Adults); and Tobacco, Alcohol, or Drug Use Assessment (Bright Futures).</p>	<p>Procedure Code(s): <i>Smoking and Tobacco Use Cessation Counseling Visit:</i> 99406, 99407</p> <p><i>Preventive Medicine, Individual Counseling:</i> 99401, 99402, 99403, 99404 Also see the codes in the Wellness Examinations row above.</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>Does not have diagnosis code requirements for the preventive benefit to apply.</p>
<p>Screening for Visual Impairment in Children</p> <p>USPSTF Rating (Sept. 2017): B The USPSTF recommends vision screening at least once in all children age 3 to 5 years to detect amblyopia or its risk factors.</p> <p>Bright Futures: Visual acuity screening is recommended for age 4 and 5 years as well as in cooperative 3 year olds. Instrument-based screening recommended for age 12 and 24 months, in addition to the well visits at 3-5 years of age.</p>	<p>Procedure Code(s): <i>Visual Acuity Screening (e.g., Snellen chart):</i> 99173</p> <p><i>Instrument-Based Screening:</i> 99174, 99177</p> <p>Diagnosis Code(s): See the Preventive Benefit Instructions.</p>	<p><i>Visual Acuity Screening (99173):</i> Up to age 21 years (ends on 22nd birthday). Does not have diagnosis code requirements for preventive benefits to apply.</p> <p><i>Instrument-Based Screening (99174 and 99177):</i></p> <ul style="list-style-type: none"> • Age 1 to 5 (ends on 6th birthday): Does not have diagnosis code requirements for preventive benefits to apply. • Age 6 to 21 years (ends on 22nd birthday): Refer to the Medical Policy titled Omnibus Codes for allowable diagnoses.
<p>Behavioral Counseling to Prevent Skin Cancer</p> <p>USPSTF Rating (March 2018): B The USPSTF recommends counseling young adults, adolescents, children, and parents of young children about minimizing exposure to ultraviolet (UV) radiation for persons ages 6 months</p>	<p>N/A</p>	<p>This service is included in a preventive care wellness examination or focused E&M visit.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
to 24 years with fair skin types to reduce their risk of skin cancer.		
<p>Falls Prevention in Community-Dwelling Older Adults</p> <p>USPSTF Rating (June 2024): B The USPSTF recommends exercise interventions to prevent falls in community-dwelling adults 65 years or older who are at increased risk for falls.</p>	N/A	This service is included in a preventive care wellness examination or focused E&M visit.
<p>Screening for Intimate Partner Violence</p> <p>USPSTF Rating (Oct. 2018): B The USPSTF recommends that clinicians screen for intimate partner violence in women of reproductive age and provide or refer women who screen positive to ongoing support services.</p> <p>Also see Screening and Counseling for Interpersonal and Domestic Violence in the <i>Expanded Women's Preventive Health</i> section.</p>	N/A	This service is included in a preventive care wellness examination.
<p>Screening for Lung Cancer with Low-Dose Computed Tomography</p> <p>USPSTF Rating (March 2021): B The USPSTF recommends annual screening for lung cancer with low-dose computed tomography (LDCT) in adults aged 50 to 80 years who have a 20 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.</p>	<p>Procedure Code(s): 71271</p> <p>Diagnosis Code(s): F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891</p> <p>Codes for Reporting Purposes: G9275, G9276</p> <p>Note: HCPCS codes G9275 and G9276, are for reporting purposes only, if applicable. These codes are not separately reimbursable.</p>	<p>Requires one of the diagnosis codes listed in this row.</p> <p>Limitations:</p> <ul style="list-style-type: none"> Limited to one per year, and All of the following criteria: <ul style="list-style-type: none"> Age 50 to 80 years (ends on 81st birthday), and At least 20 pack-years* of smoking history, and Either a current smoker or has quit within the past 15 years <p>Note: Prior authorization requirements may apply, depending on plan.</p> <p>*A pack-year is a way to measure the amount a person has smoked over a long period of time. It is calculated by multiplying the number of packs of cigarettes smoked per day by the number of years the person has smoked. For example, 1 pack year is equal to smoking 1 pack per day for 1</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
		year, or 2 packs per day for half a year, and so on. Source: National Institutes of Health, National Cancer Institute Dictionary of Cancer Terms, pack year definition web page. https://www.cancer.gov/publications/dictionaries/cancer-terms/def/pack-year
<p>Fluoride Application in Primary Care</p> <p>USPSTF Rating (May 2014): B Children From Birth Through Age 5 Years. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption.</p> <p>Bright Futures (July 2022): Bright Futures adopted the May 2014 recommendation of the USPSTF and further recommends, once teeth are present, apply fluoride varnish to all children every 3 to 6 months in the primary care or dental office, based on caries risk.</p>	<p>Procedure Code(s): <i>Application of Topical Fluoride by Physician or Other Qualified Health Care Professional:</i> 99188</p> <p>Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.</p>	<p>Age 0-5years (ends on 6th birthday).</p> <p>Does not have diagnosis code requirements for the preventive benefit to apply.</p>
<p>Latent Tuberculosis Infection in Adults: Screening</p> <p>USPSTF Rating (May 2023): B The USPSTF recommends screening for latent tuberculosis infection (LTBI) in populations at increased risk. This recommendation applies to asymptomatic adults 18 years or older at increased risk for tuberculosis (TB).</p>	<p>Procedure Code(s): <i>Screening:</i> 86480, 86481, 86580</p> <p><i>Follow-Up Visit to Check Results:</i> 99211</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): R76.11, R76.12, Z00.00, Z00.01, Z11.1, Z11.7, Z20.1</p> <p>Note for age 18-21 years (ends on 22nd birthday): In addition to the codes in this row, the preventive benefit also applies to the diagnosis codes listed in the Bright Futures row: Tuberculosis (TB) Testing.</p>	<p><i>Screening:</i> Ages 18 years and up.</p> <p>Requires one of the diagnosis codes listed in this row for CPT codes 86480, 86481, and 86580.</p> <p><i>Follow-Up Visit to Check Results (99211):</i> CPT code 99211 requires diagnosis code R76.11 or R76.12.</p> <p><i>Blood Draw:</i> Ages 18 years and up.</p> <p>Required to be billed with CPT codes 86480 or 86481 and one of the diagnosis codes listed in this row.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p><i>Hypertensive Disorders of Pregnancy – Screening</i></p> <p>USPSTF Rating (Sept. 2023): B The USPSTF recommends screening for hypertensive disorders in pregnant persons with blood pressure measurements throughout pregnancy.</p>	<p>Hypertensive disorders of pregnancy screening by blood pressure measurement is included in the code for a prenatal care office visit. See the following code groups in the <i>Expanded Women's Preventive Health</i> section:</p> <ul style="list-style-type: none"> • Prenatal Office Visits • Prenatal Care Visits • Global Obstetrical Codes 	<p>See the following code groups in the <i>Expanded Women's Preventive Health</i> section:</p> <ul style="list-style-type: none"> • Prenatal Office Visits • Prenatal Care Visits • Global Obstetrical Codes
<p><i>Perinatal Depression – Preventive Interventions (Counseling)</i></p> <p>USPSTF Rating (Feb. 2019): B The USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal depression to counseling interventions.</p> <p>Note: This policy addresses coding for interventions from a network medical provider only. For perinatal depression preventive interventions with a mental health provider, refer to the plan's mental health plan benefit administrator.</p> <p>Also see the rows for Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Depression in Children and Adolescents (Screening) (USPSTF); and Depression Screening (Bright Futures).</p>	<p>Code Group 1 Procedure Code(s): <i>Preventive Medicine Individual Counseling:</i> 99401, 99402, 99403, 99404</p> <p><i>Preventive Medicine, Group Counseling:</i> 99411, 99412</p> <p><i>Prenatal Care Visits:</i> 59425, 59426</p> <p><i>Preventive Medicine Services (Evaluation and Management):</i> 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397</p> <p>Code Group 2 Procedure Code(s): <i>Evaluation and Management (Office Visits):</i> 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, G0463</p> <p>Code Group 2 Diagnosis Code(s): A Pregnancy Diagnosis Code; or Z39.2 (encounter for routine postpartum follow-up); or Z13.32 (encounter for screening for maternal depression)</p>	<p>Code Group 1: Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>Code Group 2: Requires one of the Code Group 2 diagnosis codes listed in this row.</p>
<p><i>Prevention of Human Immunodeficiency Virus (HIV) Infection: Preexposure Prophylaxis</i></p> <p>USPSTF Rating (Aug. 2023): A The USPSTF recommends that clinicians prescribe preexposure prophylaxis using effective antiretroviral therapy to persons</p>	<p>Procedure Code(s): <i>Kidney Function Testing (Creatinine):</i> 82565, 82575</p> <p><i>Pregnancy Testing:</i> 81025, 84702, 84703</p> <p><i>Office Visits:</i> 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417,</p>	<p>Requires one of the diagnosis codes listed in this row in the primary position.</p> <p>Note: Prior authorization requirements may apply, depending on plan. Refer to the Medical Benefit Drug Policy titled Long-Acting Injectable Antiretroviral Agents for HIV.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p>who are at increased risk of HIV acquisition to decrease the risk of acquiring HIV.</p> <p>Note: This benefit also includes:</p> <ul style="list-style-type: none"> • Kidney function testing (creatinine) • Serologic testing for hepatitis B and C virus • Testing for other STIs • Pregnancy testing when appropriate • Ongoing follow-up and monitoring including HIV testing every 3 months <p>Refer to the plan's pharmacy benefit plan administrator for details on prescription medications available under the plan's preventive benefit.</p>	<p>G0463 (also see codes in the Wellness Examinations section)</p> <p><i>Antiretroviral Therapy Injection:</i> 96372 (Administration) J0739 (Injection cabotegravir, 1mg) G0012 (Administration)</p> <p><i>Counseling for PrEP to prevent HIV:</i> G0011, G0013</p> <p><i>Pharmacy Supplying Fee for HIV PrEP:</i> Q0521</p> <p>Diagnosis Code(s): Z11.3, Z11.4, Z20.2, Z20.6, Z29.81, Z72.51, Z72.52, Z72.53</p> <p>Also see the sections for:</p> <ul style="list-style-type: none"> • Behavioral Counseling to Prevent Sexually Transmitted Infections • Chlamydia Infection Screening • Gonorrhea Screening • Hepatitis B Virus Infection Screening • Hepatitis C Virus Infection Screening • HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults • Syphilis Screening 	
<h3>Bright Futures</h3>		
<p>Anemia Screening in Children (Bright Futures)</p>	<p>Procedure Code(s): <i>Anemia Screening in Children:</i> 85014, 85018</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Z00.110, Z00.111, Z00.121, Z00.129, Z13.0</p>	<p><i>Anemia Screening in Children:</i> Ages prenatal to 21 (ends on 22nd birthday). No frequency limit.</p> <p>Requires one of the diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Ages prenatal to 21 (ends on 22nd birthday).</p> <p>Required to be billed with 85014 or 85018 and one of the diagnosis codes listed in this row.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p>Hearing Tests</p> <p>Bright Futures (April 2017): <i>Hearing Tests:</i> Recommended at ages: Newborn; between 3-5 days to 2 months; 4 years; 5 years, 6 years; 8 years; 10 years; once between age 11-14 years; once between age 15-17 years; once between age 18-21 years; also recommended for those that have a positive risk assessment.</p> <p><i>Risk Assessment:</i> Recommended at ages: 4 mo, 6 mo, 9 mo, 12 mo, 15 mo, 18 mo, 24 mo, 30 mo, 3 years, 7 years, and 9 years.</p>	<p>Procedure Code(s): <i>Hearing Tests:</i> 92551, 92552, 92553, 92558, 92587, 92588, 92650, 92651, V5008</p> <p>Diagnosis Code(s): Examination of Hearing: Z01.10 Routine Child: Z00.121, Z00.129 General Exam (for 18-21years): Z00.00, Z00.01</p> <p>Note: A risk assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.</p>	<p>Ages 0-90 days: Does not have diagnosis code requirements for the preventive benefit to apply.</p> <p>Ages 91 days to 21 years (ends on 22nd birthday). Requires one of the diagnosis codes listed in this row. Limit of once per year.</p>
<p>Screening for Visual Impairment in Children (Bright Futures)</p>	<p>See row above for Screening for Visual Impairment in Children.</p>	<p>See row above Screening for Visual Impairment in Children.</p>
<p>Formal Developmental/ Autism Screening</p> <p>Bright Futures:</p> <ul style="list-style-type: none"> • A formal, standardized developmental screen is recommended during the 9 month visit. • A formal, standardized developmental screen is recommended during the 18 month visit, including a formal autism screen. • A formal, standardized autism screen is recommended during the 24 month visit. • A formal, standardized developmental screen is recommended during the 30 month visit. 	<p>Procedure Code(s): 96110</p> <p>Diagnosis Code(s): Z00.121, Z00.129, Z13.40, Z13.41, Z13.42, Z13.49</p>	<p>Ages prenatal to 2 years (ends on 3rd birthday).</p> <p>No frequency limit.</p> <p>Requires one of the diagnosis codes listed in this row.</p>
<p>Lead Screening</p> <p>Bright Futures: <i>Screening Lab Work:</i> Conduct risk assessment or screening, as appropriate, at the following intervals: 12 mo and 24 mo.</p> <p><i>Risk Assessment, and Screening if positive:</i> Recommended at 6 mo, 9</p>	<p>Procedure Code(s): <i>Lead Screening:</i> 83655</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Z00.121, Z00.129, Z77.011</p>	<p><i>Lead Screening:</i> Ages 6 months through age 6 years (ends on 7th birthday). No frequency limit.</p> <p>Requires one of the diagnosis codes listed in this row.</p> <p><i>Blood Draw:</i> Ages 6 months through age 6 years (ends on 7th birthday).</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
mo, 12 mo, 18 mo, 24 mo, 3 years, 4 years, 5 years, and 6 years.		Required to be billed with CPT code 83655 and one of the diagnosis codes in this row.
<i>Tuberculosis (TB) Testing</i> Bright Futures For age 18 years and older, also refer to the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults .	Procedure Code(s): <i>Screening:</i> 86580 <i>Follow-Up Visit to Check Results:</i> 99211 Diagnosis Code(s): R76.11, R76.12, Z20.1, Z00.121, Z00.129, Z11.1, Z11.7 Note for age 18 years and older: In addition to these codes, the preventive benefit also applies to all codes listed in the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults .	Ages prenatal to 21 (ends on 22 nd birthday). Note: For age 18 years and older, also refer to the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults No frequency limit. CPT code 86580 requires one of the diagnosis codes listed in this row. CPT code 99211 requires diagnosis code R76.11, R76.12, or Z11.1.
<i>Dyslipidemia Screening</i> Bright Futures (April 2014): <i>Risk Assessment:</i> Recommended at 24 mo, 4 years, 6 years, 8 years, 12 years, 13 years, 14 years, 15 years, 16 years. <i>Screening Lab Work:</i> Conduct if risk assessment is positive, or, at the following intervals: once between age 9-11 years; once between age 17-21 years	Procedure Code(s): <i>Dyslipidemia Screening Lab Work:</i> 80061, 82465, 83718, 83719, 83721, 83722, 84478 <i>Blood Draw:</i> 36415, 36416 Diagnosis Code(s): Z00.121, Z00.129, Z13.220 Note: A risk assessment is included in the code for a wellness examination visit; see the Wellness Examinations row above.	<i>Dyslipidemia Screening Lab Work:</i> Ages 24 months to 21 years (ends on 22 nd birthday). Requires one of the diagnosis codes listed in this row. <i>Blood Draw:</i> Ages 24 months to 21 years (ends on 22 nd birthday). Requires one of the listed Dyslipidemia Screening procedure codes listed in this row and one of the diagnosis codes listed in this row.
<i>Tobacco, Alcohol, or Drug Use Assessment</i> Bright Futures (April 2017): Bright Futures recommends tobacco, alcohol, or drug use assessment from age 11-21 years.	See codes in the rows above: <ul style="list-style-type: none"> • Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents • Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults • Unhealthy Drug Use Screening (Adults) 	See the rows above: <ul style="list-style-type: none"> • Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents • Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults • Unhealthy Drug Use Screening (Adults)

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<p><i>Behavioral/Social/Emotional Screening</i></p> <p>Bright Futures (July 2022): Bright Futures recommends behavioral/social/emotional screening annually from newborn to 21 years.</p> <p>Also see the rows for Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Perinatal Depression – Preventive Interventions (Counseling) (USPSTF); and Depression and Suicide Risk Screening (Bright Futures).</p>	<p>An assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.</p>	<p>See the Wellness Examinations row above.</p>
<p><i>Depression and Suicide Risk Screening</i></p> <p>Bright Futures (July 2022): Bright Futures recommends screening adolescents age 12-21 years for depression and suicide risk, making every effort to preserve confidentiality of the adolescent.</p> <p>Bright Futures (February 2017): <i>Maternal Depression Screening:</i> Routine screening for postpartum depression should be integrated into well-child visits at 1, 2, 4, and 6 months of age.</p> <p>Also see the rows for Screening for Anxiety (HRSA); Depression in Children and Adolescents (Screening) (USPSTF); and Perinatal Depression – Preventive Interventions (Counseling).</p>	<p>See the codes in the Depression in Children and Adolescents (Screening) row above.</p>	<p>See the Depression in Children and Adolescents (Screening) row above.</p>
<p><i>Sexually Transmitted Infections (STI)</i></p> <p>Bright Futures (April 2017): Bright Futures recommends the following: <i>STI Risk Assessment:</i> Conduct risk assessment at each of the recommended visits between 11 years – 21 years.</p>	<p>See the codes in the Chlamydia Infection Screening and Gonorrhea Screening rows above.</p>	<p>See the Chlamydia Infection Screening and Gonorrhea Screening rows above.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
<i>STI Lab Work</i> : Conduct if risk assessment is positive.		
<p>HIV Screening</p> <p>Bright Futures (April 2023): <i>HIV Risk Assessment</i>: Conduct risk assessment at age 11 years, 12 years, 13 years, 14 years, 19 years, 20 years, and 21 years.</p> <p><i>HIV Screening Lab Work</i>: Conduct at least once between age 15-21 years. Also recommended anytime between ages 11-14 years, when a risk assessment is positive. And after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently if at high risk.</p>	See the codes in the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row above.	See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row above.
<p>Sudden Cardiac Arrest (SCA) and Sudden Cardiac Death (SCD) – Risk Assessment and ECG Screening</p> <p>Bright Futures (July 2022): All children should be evaluated for conditions predisposing to SCA and SCD in the course of routine health care. A thorough and detailed history, family history, and physical examination are necessary to begin assessing SCA and SCD risk. The ECG should be the first test ordered when there is a concern for SCA risk. The ECG should be interpreted by a physician trained in recognizing electrical heart disease (i.e., a pediatric cardiologist or pediatric electrophysiologist).</p>	<p>ECG Screening for those at Risk</p> <p>Procedure Code(s): 93000, 93005, 93010</p> <p>Diagnosis Code(s): <i>Required Screening Diagnosis Codes (requires at least one):</i> Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129</p> <p><i>And requires one of the following Additional Diagnosis Codes (requires at least one):</i> I42.0, I42.1, I42.2, I45.81, I49.8, I49.9, R55, R06.00, R06.09, R53.83, R00.2, R01.0, R01.1, R03.0, Q87.40, Q87.410, Q87.418, Q87.42, Q87.43, Q87.85, Q93.52, Z82.41, Z84.81, Z82.49</p> <p>Risk Assessment A risk assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.</p>	<p><i>ECG Screening for those at Risk:</i> Limited to ages 11 years to 21 years (ends on 22nd birthday). Requires one of the Screening Diagnosis Codes listed in this row and one of the Additional Diagnosis Codes listed in this row.</p>

Preventive Care Services

Also see the [Expanded Women's Preventive Health](#) section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions
Hepatitis B Virus Infection Screening Bright Futures (July 2022): Bright Futures recommends screening between the ages 0-21 years (perform risk assessment for hepatitis B virus (HBV) infection).	See the codes in the Hepatitis B Virus Infection Screening row above.	See the Hepatitis B Virus Infection Screening row above.

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA).

For additional services covered for women, see the [Preventive Care Services](#) section above.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
Well-Woman Preventive Visits HRSA Requirement (Dec. 2021): WPSI Recommends that women receive at least one preventive care visit per year beginning in adolescence and continuing across the lifespan to ensure the provision of all recommended preventive services, including preconception and many services necessary for prenatal and interconception care, are obtained. The primary purpose of these visits should be the delivery and coordination of recommended preventive services as determined by age and risk factors. These services may be completed at a single or as part of a series of visits that take place over time to obtain all necessary services depending on a woman's age, health status, reproductive health needs, pregnancy status, and risk factors. Well-women visits also include pre-pregnancy, prenatal, postpartum and interpregnancy visits. Also see Wellness Examinations and other USPSTF	Procedure Code(s): <i>Well-Woman Visits:</i> See the Wellness Examinations row in the <i>Preventive Care Services</i> section.	<i>Well-Woman Visits:</i> See the Wellness Examinations row in the <i>Preventive Care Services</i> section.
	<i>Prenatal Office Visits:</i> Evaluation and Management (Office Visits): 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, G0463	<i>Prenatal Office Visits:</i> Requires a Pregnancy Diagnosis Code .
	<i>Pelvic Examination (add-on code):</i> 99459	<i>Pelvic Examination add-on code 99459:</i> Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.
	<i>Physician Prenatal Education, Group Setting:</i> 99078	<i>Physician Prenatal Education, Group Setting:</i> Requires a Pregnancy Diagnosis Code .
	<i>Prenatal Care (Antepartum) Visits:</i> 59425, 59426	<i>Prenatal Care (Antepartum) Visits:</i> Does not have diagnosis code requirements for the preventive benefit to apply.
	<i>Global Obstetrical Codes:</i> 59400, 59510, 59610, 59618	<i>Global Obstetrical Codes:</i> The routine, low-risk, prenatal visits portion of the code is covered as preventive.

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
recommendations during pregnancy in the <i>Preventive Care Services</i> section.	<i>Postpartum Care Visits (outpatient):</i> 59430	Does not have diagnosis code requirements for the preventive benefit to apply. <i>Postpartum Care Visits (outpatient):</i> Does not have diagnosis code requirements for the preventive benefit to apply.
<p>Screening for Diabetes in Pregnancy</p> <p>HRSA Requirement (Jan. 2023): Recommends screening pregnant women for gestational diabetes mellitus after 24 weeks of gestation (preferably between 24 and 28 weeks of gestation) to prevent adverse birth outcomes. WPSI recommends screening pregnant women with risk factors for type 2 diabetes or GDM before 24 weeks of gestation – ideally at the first prenatal visit.</p> <p>Also see the Screening for Pre-Diabetes and Type 2 Diabetes and Gestational Diabetes Screening sections of the <i>Preventive Care Services</i> section, and the Screening for Diabetes After Pregnancy section.</p>	<p>Procedure Code(s): <i>Diabetes Screening:</i> 82947, 82948, 82950, 82951, 82952, 83036</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): Pregnancy Diagnosis Codes</p>	<p><i>Diabetes Screening:</i> Requires a Pregnancy Diagnosis Code (regardless of gestational week).</p> <p><i>Blood Draw:</i> Requires one of the diabetes screening procedure codes listed in this row and one of the Pregnancy Diagnosis Codes.</p> <p>Note: If a diabetes diagnosis code is present in any position, the preventive benefit will not be applied. See the Diabetes Diagnosis Code List.</p>
<p>Screening for Diabetes After Pregnancy</p> <p>HRSA Requirement (Jan. 2023): Recommends screening for type 2 diabetes in women with a history of gestational diabetes mellitus (GDM) who are not currently pregnant and who have not previously been diagnosed with type 2 diabetes. Initial testing should ideally occur within the first year postpartum and can be conducted as early as 4-6 weeks postpartum. Women who were not screened in the first year postpartum or those with a negative initial postpartum</p>	<p>Procedure Code(s): <i>Diabetes Screening:</i> 82947, 82948, 82950, 82951, 82952, 83036</p> <p><i>Blood Draw:</i> 36415, 36416</p> <p>Diagnosis Code(s): <i>Required Screening Diagnosis Codes (requires at least one):</i> Z00.00, Z00.01, Z13.1</p> <p>And requires the following additional code: <i>Additional Diagnosis Code Required:</i> Z86.32 (personal history of gestational diabetes)</p>	<p><i>Diabetes Screening:</i> Requires one of the Required Screening diagnosis codes listed in this row and Z86.32.</p> <p>No age limit.</p> <p><i>Blood Draw:</i> Requires one of the Diabetes Screening procedure codes listed in this row and one of the Required Screening diagnosis codes listed in this row and Z86.32.</p> <p>Note: If a diabetes diagnosis code is present in any position, the preventive benefit will not be applied. See the Diabetes Diagnosis Code List.</p>

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the [Preventive Care Services](#) section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
<p>screening test result should be screened at least every 3 years for a minimum of 10 years after pregnancy. For those with a positive screening test result in the early postpartum period, testing should be repeated at least 6 months postpartum to confirm the diagnosis of diabetes regardless of the type of initial test (e.g., fasting plasma glucose, hemoglobin A1c, in the first 6 months postpartum regardless of whether the test results are positive or negative because the hemoglobin A1c test is less accurate during the first 6 months postpartum.</p> <p>Also see Gestational Diabetes Screening and Screening for Pre-Diabetes and Type 2 in the <i>Preventive Care Services</i> section, and the Screening for Diabetes in Pregnancy section.</p>		
<p>Screening for Urinary Incontinence</p> <p>HRSA Requirement (Jan. 2024) The Women's Preventive Services Initiative recommends screening women for urinary incontinence annually. Screening should address whether women experience urinary incontinence and whether it impacts their activities and quality of life. If indicated, facilitating further evaluation and treatment is recommended.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>
<p>Counseling for Sexually Transmitted Infections (STIs)</p> <p>HRSA Requirement (Dec. 2021): WPSI recommends directed behavioral counseling by a health care clinician or other appropriately trained individual for sexually active adolescent and adult women at an increased risk for STIs. WPSI recommends that clinicians review</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the [Preventive Care Services](#) section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
<p>a woman's sexual history and risk factors to help identify those at an increased risk of STIs. Risk factors include, but are not limited to, age younger than 25, a recent history of an STI, a new sex partner, multiple partners, a partner with concurrent partners, a partner with an STI, and a lack of or inconsistent condom use.</p> <p>For adolescents and women not identified as high risk, counseling to reduce the risk of STIs should be considered, as determined by clinical judgment.</p>		
<p>Screening for Human Immunodeficiency Virus Infection (HIV)</p> <p>HRSA Requirement (Dec. 2021): The Women's Preventive Services Initiative (WPSI) recommends all adolescent and adult women, ages 15 and older, receive a screening test for human immunodeficiency virus (HIV) at least once during their lifetime. Earlier or additional screening should be based on risk, and rescreening annually or more often may be appropriate beginning at age 13 for adolescent and adult women with an increased risk of HIV infection. The WPSI recommends risk assessment and prevention education for HIV infection beginning at age 13 and continuing as determined by risk. A screening test for HIV is recommended for all pregnant women upon initiation of prenatal care with rescreening during pregnancy based on risk factors. Rapid HIV testing is recommended for pregnant women who present in labor with an undocumented HIV status.</p>	<p>Education and Risk Assessment</p> <p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above</p> <p>Screening Tests</p> <p>See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row in the <i>Preventive Care Services</i> section above</p>	<p>Education and Risk Assessment</p> <p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p> <p>Screening Tests</p> <p>See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row in the <i>Preventive Care Services</i> section above.</p>

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
<p>Contraceptive Methods (Including Sterilizations)</p> <p>HRSA Requirement (Dec. 2021): WPSI recommends that adolescent and adult women have access to the full range of contraceptives and contraceptive care to prevent unintended pregnancies and improve birth outcomes. Contraceptive care includes screening, education, counseling, and provision of contraceptives (including in the postpartum period). Contraceptive care also includes follow-up care (e.g., management, evaluation, and changes, including the removal, continuation, and discontinuation of contraceptives). WPSI recommends that the full range of U.S. Food and Drug Administration (FDA)-approved, -granted, or -cleared contraceptives, effective family planning practices, and sterilization procedures be available as part of contraceptive care. The full range of contraceptives includes those currently listed in the FDA's Birth Control Guide: (1) sterilization surgery for women, (2) implantable rods, (3) copper intrauterine devices, (4) intrauterine devices with progestin (all durations and doses), (5) injectable contraceptives, (6) oral contraceptives (combined pill), (7) oral contraceptives (progestin only), (8) oral contraceptives (extended or continuous use), (9) the contraceptive patch, (10) vaginal contraceptive rings, (11) diaphragms, (12) contraceptive sponges, (13) cervical caps, (14) condoms, (15) spermicides, (16) emergency contraception</p>	<p>Code Group 1 Procedure Code(s): <i>Sterilizations:</i> Tubal Ligation, Oviduct Occlusion: 58600, 58605, 58611, 58615, 58670, 58671, A4264 (See Code Group 4 below for Tubal Ligation Follow-Up) <i>Contraceptive Methods:</i> Diaphragm or Cervical Cap: 57170, A4261, A4266 IUD (copper): J7300 IUD (Skyla®): J7301 IUD (Liletta®): J7297 IUD (Kyleena®): J7296 (See Code Group 2 below for additional IUD codes)</p>	<p>Code Group 1: Does not have diagnosis code requirements for preventive benefits to apply.</p>
	<p>Code Group 2 Procedure Code(s): <i>Contraceptive Methods:</i> Implantable Devices: J7306, J7307 11976 (capsule removal) 11981 (implant insertion) 11982 (implant removal) 11983 (removal with reinsertion) IUDs: J7298 (Mirena®) S4989 58300, S4981 (insertion) 58301 (removal) (See Code Group 1 above for additional IUD codes) <i>Injections:</i> 96372 (administration) J1050 (injection)</p> <p>Code Group 2 Diagnosis Code(s): These are required for Code Group 2. <i>Contraceptive Management:</i> Z30.012, Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.09, Z30.40, Z30.42, Z30.430, Z30.431, Z30.432, Z30.433, Z30.46, Z30.49, Z30.8, Z30.9</p>	<p>Code Group 2: Requires one of the Code Group 2 diagnosis codes listed in this row.</p>

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the [Preventive Care Services](#) section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
<p>(levonorgestrel), and (17) emergency contraception (ulipristal acetate), and any additional contraceptives approved, granted, or cleared by the FDA.</p> <p>Notes:</p> <ul style="list-style-type: none"> Coverage includes member reimbursement for the cost of FDA-approved, cleared, or granted mobile device applications for use as contraception consistent with the FDA-approved, cleared, or granted indication. For counseling and follow-up care, see the Wellness Examinations row in the <i>Preventive Care Services</i> section above. Certain employers may qualify for an exemption from covering contraceptive methods and sterilizations on account of religious objections. Refer to the plan's pharmacy benefit plan administrator for details on pharmacy contraceptives available under the plan's preventive benefit. <p>Also refer to the Medical Policy titled Outpatient Surgical Procedures – Site of Service.</p>	<p>Code Group 3 Procedure Code(s): <i>Anesthesia for Sterilization:</i> 00851, 00940, 00942, 00950, 00952, 01960, 01961, 01965, 01966, 01967, 01968</p> <p><i>Sterilization - Laparoscopy with Removal of Adnexal Structures:</i> 58661</p> <p>Code Group 3 Diagnosis Code(s): This code is required for all Code Group 3 Procedure Codes: <i>Sterilization:</i> Z30.2</p> <p>Tubal Ligation Follow-Up Hysterosalpingogram</p> <p>Code Group 4 Procedure Code(s): <i>Catheterization and Introduction of Saline or Contrast Material:</i> 58340</p> <p><i>Hysterosalpingography:</i> 74740</p> <p><i>Contrast Material:</i> Q9967</p> <p>Code Group 4 Diagnosis Code(s): <i>Tubal Ligation Status:</i> Z98.51</p>	<p>Code Group 3: Requires one of the Code Group 3 diagnosis code listed in this row.</p> <p>Code Group 4: Requires one of the Code Group 4 diagnosis code listed in this row.</p>
	<p>Code Group 5 Procedure Code(s): <i>IUD Follow-Up Evaluation and Management (Office Visit):</i> 99211, 99212</p> <p><i>Pelvic Examination (add-on code):</i> 99459</p> <p>Refer to Code Group 7, Related Visits section below, for additional coding for Evaluation and Management (Office Visits).</p> <p>Code Group 5 Diagnosis Code(s): <i>Encounter for routine checking of intrauterine contraceptive device:</i> Z30.431</p>	<p>Code Group 5: Requires one of the Code Group 5 diagnosis code listed in this row.</p> <p><i>Pelvic Examination add-on code 99459:</i> Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.</p>

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the [Preventive Care Services](#) section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
	<p>Code Group 6 Procedure Code(s): <i>Impacted IUD removal</i> 58562</p> <p>Code Group 6 Diagnosis Code(s): Z30.432, Z30.433</p> <p>Code Group 7 Procedure Code(s): <i>Related Visits: Related Evaluation and Management Office/Outpatient Visits for Contraception or Sterilization:</i> 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, G0463</p> <p><i>Pelvic Examination (add-on code):</i> 99459 Also see coding in the Wellness Examinations row above.</p> <p><i>Related Pregnancy Tests:</i> Pregnancy Tests When Related to Contraception or Sterilization: 81025, 84702, 84703</p> <p>Code Group 7 Diagnosis Codes: <i>Tubal Ligation Status:</i> Z98.51 <i>Sterilization:</i> Z30.2</p> <p><i>Contraceptive Management:</i> Z30.012, Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.09, Z30.40, Z30.42, Z30.430, Z30.431, Z30.432, Z30.433, Z30.46, Z30.49, Z30.8, Z30.9</p>	<p>Code Group 6: Requires one of the Code Group 6 diagnosis codes listed in this row.</p> <p>Code Group 7: Requires one of the Code Group 7 diagnosis codes listed in this row.</p> <p><i>Pelvic Examination add-on CPT code 99459:</i> Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.</p>
<p>Breastfeeding Services and Supplies</p> <p>HRSA Requirement (Dec. 2021): WPSI recommends comprehensive lactation support services including consultation; counseling; education by clinicians and peer support services; and breastfeeding equipment and supplies) during the antenatal, perinatal, and postpartum periods to optimize the successful initiation and maintenance of breastfeeding. Breastfeeding equipment and supplies include, but are not limited</p>	<p>Counseling and Education Procedure Code(s): 98960, 98961, 98962, 99242*, 99243*, 99244*, 99245*, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, S9443</p> <p>Also see the codes in the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p> <p>Diagnosis Code(s): B37.89, N61.1, N64.4, N64.51, N64.52, N64.53, N64.59, N64.89, O91.011, O91.012, O91.013, O91.019, O91.02, O91.03, O91.111, O91.112, O91.113, O91.119, O91.13,</p>	<p>Counseling and Education</p> <p>Requires one of the diagnosis codes listed in this row for CPT codes 98960-98962, 99242-99245, 99341-99345, and 99347-99350.</p> <p>Does not have diagnosis code requirements for preventive benefits to apply for HCPCS code S9443.</p>

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the [Preventive Care Services](#) section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
<p>to, double electric breast pumps (including pump parts and maintenance) and breast milk storage supplies. Access to double electric pumps should be a priority to optimize breastfeeding and should not be predicated on prior failure of a manual pump. Breastfeeding equipment may also include equipment and supplies as clinically indicated to support dyads with breastfeeding difficulties and those who need additional services.</p>	<p>O91.211, O91.212, O91.213, O91.219, O91.22, O91.23, O92.011, O92.012, O92.013, O92.019, O92.02, O92.03, O92.111, O92.112, O92.113, O92.119, O92.12, O92.13, O92.20, O92.29, O92.3, O92.4, O92.5, O92.70, O92.79, Q83.1, Q83.2, Q83.3, Q83.8, Z39.1, Z39.2</p> <p>*For additional information on the reimbursement of consultation codes 99242-99245, refer to the Reimbursement Policy titled Consultation Services.</p> <p>Breastfeeding Equipment & Supplies Procedure Code(s): <i>Personal Use Electric Breast Pump:</i> E0603</p> <p><i>Breast Pump Supplies:</i> A4281, A4282, A4283, A4284, A4285, A4286, A4287</p> <p>Diagnosis Code(s): Pregnancy Diagnosis Codes or Z39.1.</p>	<p>Breastfeeding Equipment & Supplies HCPCS code E0603 is limited to one purchase per birth.</p> <p>HCPCS codes A4281-A4287 and E0603 require at least one of the diagnosis codes listed in this row.</p>
<p>Screening and Counseling for Interpersonal and Domestic Violence</p> <p>HRSA Requirement (Dec. 2016): Recommends screening adolescents and women for interpersonal and domestic violence, at least annually, and, when needed, providing or referring for initial intervention services. Interpersonal and domestic violence includes physical violence, sexual violence, stalking and psychological aggression (including coercion), reproductive coercion, neglect, and the threat of violence, abuse, or both. Intervention services include, but are not limited to, counseling, education, harm reduction strategies, and referral to appropriate supportive services.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA).
For additional services covered for women, see the [Preventive Care Services](#) section above.
Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
Also see the Screening for Intimate Partner Violence row in the <i>Preventive Care Services</i> section above.		
<p><i>Breast Cancer Screening for Average-Risk Women</i></p> <p>HRSA Requirement (Dec. 2016): Recommends that average-risk women initiate mammography screening no earlier than age 40 and no later than age 50. Screening mammography should occur at least biennially and as frequently as annually. Screening should continue through at least age 74 and age alone should not be the basis to discontinue screening. These screening recommendations are for women at average risk of breast cancer. Women at increased risk should also undergo periodic mammography screening; however, recommendations for additional services are beyond the scope of this recommendation.</p>	See the Screening Mammography row in the <i>Preventive Care Services</i> section above.	See the Screening Mammography row in the <i>Preventive Care Services</i> section above.
<p><i>Screening for Cervical Cancer</i></p> <p>HRSA Requirement (Dec. 2016): Recommends cervical cancer screening for average-risk women aged 21 to 65 years. For women aged 21 to 29 years recommends cervical cancer screening using cervical cytology (Pap test) every 3 years. Co-testing with cytology and human papillomavirus testing is not recommended for women younger than 30 years. Women aged 30 to 65 years should be screened with cytology and human papillomavirus testing every 5 years or cytology alone every 3 years. Women who are at average risk should not be screened more than once every 3 years.</p>	<p><i>Human Papillomavirus DNA Testing (HPV)</i> See the Cervical Cancer Screening row in the <i>Preventive Care Services</i> section above.</p> <p><i>Cervical Cytology (Pap Test)</i> See the Cervical Cancer Screening row in the <i>Preventive Care Services</i> section above.</p>	<p><i>Human Papillomavirus DNA Testing (HPV)</i> See the Cervical Cancer Screening row in the <i>Preventive Care Services</i> section above.</p> <p><i>Cervical Cytology (Pap Test)</i> See the Cervical Cancer Screening row in the <i>Preventive Care Services</i> section above.</p>

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the [Preventive Care Services](#) section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions
<p>Screening for Anxiety</p> <p>HRSA Requirement (Dec. 2019): The Women's Preventive Services Initiative recommends screening for anxiety in adolescent and adult women, including those who are pregnant or postpartum. Optimal screening intervals are unknown and clinical judgement should be used to determine screening frequency. Given the high prevalence of anxiety disorders, lack of recognition in clinical practices, and multiple problems associated with untreated anxiety, clinicians should consider screening women who have not been recently screened.</p> <p>Also see the rows for Anxiety Disorders in Adults Screening (USPSTF); Screening for Anxiety in Children and Adolescents (USPSTF); Screening for Depression in Adults (USPSTF); Depression in Children and Adolescents (Screening) (USPSTF); Perinatal Depression – Preventive Interventions (Counseling) (USPSTF); and Depression Screening (Bright Futures) in the <i>Preventive Care Services</i> section above.</p>	<p>Procedure Code(s): 96127</p> <p>Diagnosis Code(s): <i>Encounter for Screening Examination for Other Mental Health and Behavioral Disorders:</i> Z13.39</p>	<p>Requires the diagnosis code listed in this row.</p>
<p>Obesity Prevention in Midlife Women (Counseling)</p> <p>HRSA Requirement (Dec. 2021): WPSI recommends counseling midlife women aged 40 to 60 years with normal or overweight body mass index (BMI) (18.5-29.9 km/m²) to maintain weight or limit weight gain to prevent obesity. Counseling may include individualized discussion of healthy eating and physical activity.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>	<p>See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.</p>

Diagnosis Codes

[Preventive Care Services: ICD-10 Diagnosis Codes](#)

Preventive Vaccine Codes

Preventive Care Services: Vaccine Codes

Benefit Considerations

Certain plans are not required to include coverage for the services identified by the federal Patient Protection and Affordable Care Act (PPACA). Refer to the member specific benefit plan document for coverage details.

Clinical Evidence

Refer to the *Service* column in the [Applicable Codes](#) section for the recommendation statements supporting this policy.

References

ACIP Vaccine-Specific Recommendations: https://www.cdc.gov/acip-recs/hcp/vaccine-specific/?CDC_AAref_Val=https://www.cdc.gov/vaccines/hcp/acip-recs/index.html. Accessed December 9, 2024.

American Academy of Family Physicians (AAFP) Clinical Preventive Services Recommendations: <https://www.aafp.org/family-physician/patient-care/clinical-recommendations/clinical-practice-guidelines/clinical-preventive-services-recommendations.html>. Accessed December 9, 2024.

American Academy of Pediatrics, Bright Futures Guidelines, 4th edition, Evidence and Rationale chapter: <https://www.aap.org/en/practice-management/bright-futures/bright-futures-materials-and-tools/bright-futures-guidelines-and-pocket-guide/>. Accessed December 9, 2024.

American Academy of Pediatrics/Bright Futures/Recommendations for Pediatric Preventive Healthcare. (For ages 0-21): https://downloads.aap.org/AAP/PDF/periodicity_schedule.pdf. Accessed December 9, 2024.

American Academy of Pediatrics: <http://www.aap.org/>. Accessed December 9, 2024.

Centers for Disease Control and Prevention/Immunization Schedules: <https://www.cdc.gov/vaccines/hcp/imz-schedules/index.html>. Accessed December 9, 2024.

Grade Definitions for USPSTF Recommendations: <http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions>. Accessed December 9, 2024.

July 19, 2010 IRS Interim Rules: http://www.irs.gov/irb/2010-29_IRB/index.html. Accessed December 9, 2024.

Published Recommendations, U.S. Preventive Services Task Force: <http://www.uspreventiveservicestaskforce.org/BrowseRec/Index/browse-recommendations>. Accessed December 9, 2024.

U.S. Food and Drug Administration (FDA), Vaccines Licensed for Use in the United States: <http://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/UCM093833>. Accessed December 9, 2024.

U.S. Vaccine Names Information Page: <https://www.cdc.gov/vaccines/hcp/vaccines-us/index.html>. Accessed December 9, 2024.

Women's Preventive Services Guidelines (HRSA): <https://www.hrsa.gov/womens-guidelines>. Accessed December 9, 2024.

Women's Preventive Services Initiative (WPSI): <https://www.womenspreventivehealth.org/recommendations/>. Accessed December 9, 2024.

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

This Medical Policy may also be applied to Medicare Advantage plans in certain instances. In the absence of a Medicare National Coverage Determination (NCD), Local Coverage Determination (LCD), or other Medicare coverage guidance,

CMS allows a Medicare Advantage Organization (MAO) to create its own coverage determinations, using objective evidence-based rationale relying on authoritative evidence ([Medicare IOM Pub. No. 100-16, Ch. 4, §90.5](#)).

UnitedHealthcare may also use tools developed by third parties, such as the InterQual[®] criteria, to assist us in administering health benefits. UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.